

ABNORMALITIES OF THE UPPER LIMB

ABNORMALITIES OF THE UPPER LIMB

Minor finger abnormalities are common (Table 44.11 not all require surgical intervention. Comfort and function are more important than appearance. Robert Kienböck , 1871-1953, Professor of Radiology , Vienna, Austria, described this condition in 1910. Hans Jessen Panner , 1871-1930, radiologist, Copenhagen, Denmark, described this condition in 1927. Albert Henry Freiberg , 1869-1940, Professor of Orthopaedic Surgery , The University of Cincinnati, Cincinnati, OH, USA, described this disease in 1926. Alban Köhler , 1874-1947, radiologist, of Wiesbaden, Germany , described this disease in 1908. James Warren Sever , 1878-1964, orthopaedic surgeon, The Children's Hospital, Boston, MA, USA, described apophysitis of the os calcis in 1912. - - - - Function is also the most important consideration when managing more extensive upper limb abnormalities. Treatment is often delayed until hand dominance is established and it is clear what problems a specific deformity is causing any given child. Children are very adaptable and cope with disability readily than their parents/doctors expect. abilities much more

(b) Figure 44.31 Congenital vertical talus: (a) lateral photograph demonstrating

strating a 'rocker-bottom foot'; (b) lateral radiograph showing hindfoot equinus and suggesting dorsal subluxation of the non-ossi

/f_i ed navic

ular and forefoot with respect to the head of the talus.

ABNORMALITIES OF THE UPPER LIMB

Minor finger abnormalities are common (Table 44.11 not all require surgical intervention. Comfort and function are more important than appearance. Robert Kienböck , 1871-1953, Professor of Radiology , Vienna, Austria, described this condition in 1910. Hans Jessen Panner , 1871-1930, radiologist, Copenhagen, Denmark, described this condition in 1927. Albert Henry Freiberg , 1869-1940, Professor of Orthopaedic Surgery , The University of Cincinnati, Cincinnati, OH, USA, described this disease in 1926. Alban Köhler , 1874-1947, radiologist, of Wiesbaden, Germany , described this disease in 1908. James Warren Sever , 1878-1964, orthopaedic surgeon, The Children's Hospital, Boston, MA, USA, described apophysitis of the os calcis in 1912. - - - - Function is also the most important consideration when managing more extensive upper limb abnormalities. Treatment is often delayed until hand dominance is established and it is clear what problems a specific deformity is causing any given child. Children are very adaptable and cope with dis - e readily than their parents/doctors expect. abilities much mor

(b) Figure 44.31 Congenital vertical talus: (a) lateral photograph demon

strating a 'rocker-bottom foot'; (b) lateral radiograph showing hindfoot equinus and suggesting dorsal subluxation of the non-ossi /f_i ed navic

ular and forefoot with respect to the head of the talus.

ABNORMALITIES OF THE UPPER LIMB

Minor finger abnormalities are common (Table 44.11 not all require surgical intervention. Comfort and function are more important than appearance. Robert Kienböck , 1871–1953, Professor of Radiology , Vienna, Austria, described this condition in 1910. Hans Jessen Panner , 1871–1930, radiologist, Copenhagen, Denmark, described this condition in 1927. Albert Henry Freiberg , 1869–1940, Professor of Orthopaedic Surgery , The University of Cincinnati, Cincinnati, OH, USA, described this disease in 1926. Alban Köhler , 1874–1947, radiologist, of Wiesbaden, Germany , described this disease in 1908. James Warren Sever , 1878–1964, orthopaedic surgeon, The Children’s Hospital, Boston, MA, USA, described apophysitis of the os calcis in 1912. - - - Function is also the most important consideration when managing more extensive upper limb abnormalities. Treatment is often delayed until hand dominance is established and it is clear what problems a specific deformity is causing any given child. Children are very adaptable and cope with disability much more readily than their parents/doctors expect. abilities much more

(b) Figure 44.31 Congenital vertical talus: (a) lateral photograph demonstrating

strating a ‘rocker-bottom foot’; (b) lateral radiograph showing hindfoot equinus and suggesting dorsal subluxation of the non-ossified navicular

ular and forefoot with respect to the head of the talus.