

Ankylosing spondylitis

Ankylosing spondylitis

Should a patient with ankylosing spondylitis present following trauma, a high index of suspicion for occult fractures should be present. It is common for patients with ankylosing spondylitis to develop epidural haematomas with subtle neurological deficit. Patients with a significant fixed flexion deformity at the cervicothoracic junction ('chin-on-chest' deformity), limited forward gaze and eating and swallowing difficulties may be treated with a closing wedge osteotomy at the cervicothoracic junction (Figure 37.7). Extension osteotomies can also be performed in the thoracic and lumbar spine. Ankylosing spondylitis

Should a patient with ankylosing spondylitis present following trauma, a high index of suspicion for occult fractures should be present. It is common for patients with ankylosing spondylitis to develop epidural haematomas with subtle neurological deficit. Patients with a significant fixed flexion deformity at the cervicothoracic junction ('chin-on-chest' deformity), limited forward gaze and eating and swallowing difficulties may be treated with a closing wedge osteotomy at the cervicothoracic junction (Figure 37.7). Extension osteotomies can also be performed in the thoracic and lumbar spine. Ankylosing spondylitis

Should a patient with ankylosing spondylitis present following trauma, a high index of suspicion for occult fractures should be present. It is common for patients with ankylosing spondylitis to develop epidural haematomas with subtle neurological deficit. Patients with a significant fixed flexion deformity at the cervicothoracic junction ('chin-on-chest' deformity), limited forward gaze and eating and swallowing difficulties may be treated with a closing wedge osteotomy at the cervicothoracic junction (Figure 37.7). Extension osteotomies can also be performed in the thoracic and lumbar spine.

Revision #1

Created 2025-12-31 15:15:06 UTC by Omar Ayman

Updated 2025-12-31 15:15:06 UTC by Omar Ayman