

ANORECTAL PROBLEMS

Constipation

ANORECTAL PROBLEMS Constipation

The passage of hard or infrequent stools may be secondary to an anal fissure, Hirschsprung's disease, an anorectal malformation, a neuropathic bowel, a mega-rectosigmoid or idiopathic constipation. A detailed history and examination of the abdomen, anus and spine identify most causes. A rectal biopsy may be needed to exclude late presenting Hirschsprung's disease. In the absence of surgical pathology, the child is best managed by a paediatrician. ANORECTAL PROBLEMS Constipation

The passage of hard or infrequent stools may be secondary to an anal fissure, Hirschsprung's disease, an anorectal malformation, a neuropathic bowel, a mega-rectosigmoid or idiopathic constipation. A detailed history and examination of the abdomen, anus and spine identify most causes. A rectal biopsy may be needed to exclude late presenting Hirschsprung's disease. In the absence of surgical pathology, the child is best managed by a paediatrician. ANORECTAL PROBLEMS Constipation

The passage of hard or infrequent stools may be secondary to an anal fissure, Hirschsprung's disease, an anorectal malformation, a neuropathic bowel, a mega-rectosigmoid or idiopathic constipation. A detailed history and examination of the abdomen, anus and spine identify most causes. A rectal biopsy may be needed to exclude late presenting Hirschsprung's disease. In the absence of surgical pathology, the child is best managed by a paediatrician.

Revision #1

Created 2025-12-31 15:18:24 UTC by Omar Ayman

Updated 2025-12-31 15:18:24 UTC by Omar Ayman