

# Antenatal fetal hydronephrosis

## Antenatal fetal hydronephrosis

Hydronephrosis is a dilated renal pelvis that is found in 1% of antenatal scans and most commonly it resolves, especially if the dilatation is mild to moderate. Severe dilatation is associated with urinary tract obstruction or vesicoureteral reflux. Antenatal interventions are rarely indicated except in posterior urethral valves (PUVs), but postnatal imaging is needed to confirm resolution or make a diagnosis. Amniotic fluid is principally fetal urine; thus, if there is antenatal bilateral hydronephrosis with decreased amniotic fluid, PUV is a likely cause. If the PUV obstruction is thought to be damaging the kidneys, then there may be a role for an antenatal vesicoamniotic shunt to take the pressure off the upper tracts. Antenatal fetal hydronephrosis

Hydronephrosis is a dilated renal pelvis that is found in 1% of antenatal scans and most commonly it resolves, especially if the dilatation is mild to moderate. Severe dilatation is associated with urinary tract obstruction or vesicoureteral reflux. Antenatal interventions are rarely indicated except in posterior urethral valves (PUVs), but postnatal imaging is needed to confirm resolution or make a diagnosis. Amniotic fluid is principally fetal urine; thus, if there is antenatal bilateral hydronephrosis with decreased amniotic fluid, PUV is a likely cause. If the PUV obstruction is thought to be damaging the kidneys, then there may be a role for an antenatal vesicoamniotic shunt to take the pressure off the upper tracts. Antenatal fetal hydronephrosis

Hydronephrosis is a dilated renal pelvis that is found in 1% of antenatal scans and most commonly it resolves, especially if the dilatation is mild to moderate. Severe dilatation is associated with urinary tract obstruction or vesicoureteral reflux. Antenatal interventions are rarely indicated except in posterior urethral valves (PUVs), but postnatal imaging is needed to confirm resolution or make a diagnosis. Amniotic fluid is principally fetal urine; thus, if there is antenatal bilateral hydronephrosis with decreased amniotic fluid, PUV is a likely cause. If the PUV obstruction is thought to be damaging the kidneys, then there may be a role for an antenatal vesicoamniotic shunt to take the pressure off the upper tracts.

---

Revision #1

Created 2025-12-31 15:10:22 UTC by Omar Ayman

Updated 2025-12-31 15:10:22 UTC by Omar Ayman