

ASSESSMENT Light microscopy

ASSESSMENT Light microscopy

Most tissue assessment depends on conventional light microscopy. Microscopes have several lenses with various powers of magnification, typically ranging from $\times 20$ to $\times 400$ or more. A low-power lens allows scanning of a sample and assessment while a higher power lens allows a view of overall architecture, with closer view with more detail (Figure 11.21). Attachment of one or more teaching arms, a camera and other accessories are possible for many microscopes (Figure 11.8). Polarisation assists with detection of some types of foreign material (e.g. sutures) or to assess a special stain (e.g. Congo red for amyloid deposition).

(b) Figure 11.21 (a) Low-power view of an umbilical nodule. Glands are distributed irregularly through the tissue. (b) High-power view shows benign columnar epithelium lining the glands (arrow), indicating endometriosis rather than carcinoma.

ASSESSMENT Light microscopy

Most tissue assessment depends on conventional light microscopy. Microscopes have several lenses with various powers of magnification, typically ranging from $\times 20$ to $\times 400$ or more. A low-power lens allows scanning of a sample and assessment while a higher power lens allows a view of overall architecture, with closer view with more detail (Figure 11.21). Attachment of one or more teaching arms, a camera and other accessories are possible for many microscopes (Figure 11.8). Polarisation assists with detection of some types of foreign material (e.g. sutures) or to assess a special stain (e.g. Congo red for amyloid deposition).

(b) Figure 11.21 (a) Low-power view of an umbilical nodule. Glands are distributed irregularly through the tissue. (b) High-power view shows benign columnar epithelium lining the glands (arrow), indicating endometriosis rather than carcinoma.

ASSESSMENT Light microscopy

Most tissue assessment depends on conventional light microscopy. Microscopes have several lenses with various powers of magnification, typically ranging from $\times 20$ to $\times 400$ or more. A low-power lens allows scanning of a sample and assessment while a higher power lens allows a view of overall architecture, with closer view with more detail (Figure 11.21). Attachment of one or more teaching arms, a camera and other accessories are possible for many microscopes (Figure 11.8). Polarisation assists with detection of some types of foreign material (e.g. sutures) or to assess a special stain (e.g. Congo red for amyloid deposition).

(b) Figure 11.21 (a) Low-power view of an umbilical nodule. Glands are distributed irregularly through the tissue. (b) High-power view shows benign columnar epithelium lining the glands (arrow), indicating endometriosis rather than carcinoma.

Revision #1

Created 2025-12-31 15:08:16 UTC by Omar Ayman

Updated 2025-12-31 15:08:16 UTC by Omar Ayman