

Basic methods in diagnostic molecular pathology

Basic methods in diagnostic molecular pathology

In situ hybridisation H. In situ hybridisation (ISH) uses a labelled oligonucleotide probe - that targets a specific sequence of RNA or DNA. It allows visu - alisation of the presence or absence and location of a particular RNA or DNA sequence in situ in tissue sections. Visualisation may depend on autoradiography , fluorescence microscopy - or bright-field microscopy . Chromogenic in situ hybridisation (CISH) combines ISH and immunohistochemistry for the detection of specific nucleic acid sequences and is a common alternative to fluorescence in situ hybridisation (FISH) for the detection of HER2 amplification. Viral genomes, e.g. EBV - (Figure 11.28), CMV and high-risk HPV types are detectable using this approach. ISH plays an important role in tissue diagnostics and the management of tumours.

Figure 11.28 In situ hybridisation for Epstein-Barr virus (EBV) showing extensive nuclear positivity (black nuclei) in an EBV-positive gastric adenocarcinoma.

Basic methods in diagnostic molecular pathology

In situ hybridisation H. In situ hybridisation (ISH) uses a labelled oligonucleotide probe - that targets a specific sequence of RNA or DNA. It allows visu - alisation of the presence or absence and location of a particular RNA or DNA sequence in situ in tissue sections. Visualisation may depend on autoradiography , fluorescence microscopy - or bright-field microscopy . Chromogenic in situ hybridisation (CISH) combines ISH and immunohistochemistry for the detection of specific nucleic acid sequences and is a common alternative to fluorescence in situ hybridisation (FISH) for the detection of HER2 amplification. Viral genomes, e.g. EBV - (Figure 11.28), CMV and high-risk HPV types are detectable using this approach. ISH plays an important role in tissue diagnostics and the management of tumours.

Figure 11.28 In situ hybridisation for Epstein-Barr virus (EBV) showing extensive nuclear positivity (black nuclei) in an EBV-positive gastric adenocarcinoma.

Basic methods in diagnostic molecular pathology

In situ hybridisation H. In situ hybridisation (ISH) uses a labelled oligonucleotide probe - that targets a specific sequence of RNA or DNA. It allows visu - alisation of the presence or absence and location of a particular RNA or DNA sequence in situ in tissue sections. Visualisation may depend on autoradiography , fluorescence microscopy - or bright-field microscopy . Chromogenic in situ hybridisation (CISH) combines ISH and immunohistochemistry for the detection of specific nucleic

acid sequences and is a common alternative to fluorescence in situ hybridisation (FISH) for the detection of HER2 amplification. Viral genomes, e.g. EBV - (Figure 11.28), CMV and high-risk HPV types are detectable using this approach. ISH plays an important role in tissue diagnostics and the management of tumours.

Figure 11.28 In situ hybridisation for Epstein-Barr virus (EBV) showing extensive nuclear positivity (black nuclei) in an EBV-positive gastric adenocarcinoma.

Revision #1

Created 2025-12-31 15:08:17 UTC by Omar Ayman

Updated 2025-12-31 15:08:17 UTC by Omar Ayman