

Colon

Colon

Blunt injuries to the colon are relatively infrequent; penetrating injuries occur more often. If relatively little contamination is present and the viability is satisfactory, such wounds can be repaired primarily. If, however, there is extensive contamination, the patient is physiologically compromised or the bowel is of doubtful viability, then the bowel can be closed off ('clip and drop'). A defunctioning colostomy can be formed later or the bowel reanastomosed once the patient is stable. Colon

Blunt injuries to the colon are relatively infrequent; penetrating injuries occur more often. If relatively little contamination is present and the viability is satisfactory, such wounds can be repaired primarily. If, however, there is extensive contamination, the patient is physiologically compromised or the bowel is of doubtful viability, then the bowel can be closed off ('clip and drop'). A defunctioning colostomy can be formed later or the bowel reanastomosed once the patient is stable. Colon

Blunt injuries to the colon are relatively infrequent; penetrating injuries occur more often. If relatively little contamination is present and the viability is satisfactory, such wounds can be repaired primarily. If, however, there is extensive contamination, the patient is physiologically compromised or the bowel is of doubtful viability, then the bowel can be closed off ('clip and drop'). A defunctioning colostomy can be formed later or the bowel reanastomosed once the patient is stable.

Revision #1

Created 2025-12-31 15:12:29 UTC by Omar Ayman

Updated 2025-12-31 15:12:29 UTC by Omar Ayman