

# Computed tomography

## Computed tomography

This investigation is the best test for assessing bone anatomy . Three-dimensional reconstructions are often useful for the assessment of congenital spinal deformity . However, one should remember that a typical CT of the lumbar spine will expose the patient to an effective dose of 5–10 millisieverts (mSv), which would be equivalent to 2.5–5 years of natural background radiation; those who travel on 7-hour flights (0.05 /uni00A0 mSv per 7-hour flight) would need to make 100–200 journeys in their lifetime to be exposed to the same effective dose. Computed tomography

This investigation is the best test for assessing bone anatomy . Three-dimensional reconstructions are often useful for the assessment of congenital spinal deformity . However, one should remember that a typical CT of the lumbar spine will expose the patient to an effective dose of 5–10 millisieverts (mSv), which would be equivalent to 2.5–5 years of natural background radiation; those who travel on 7-hour flights (0.05 /uni00A0 mSv per 7-hour flight) would need to make 100–200 journeys in their lifetime to be exposed to the same effective dose. Computed tomography

This investigation is the best test for assessing bone anatomy . Three-dimensional reconstructions are often useful for the assessment of congenital spinal deformity . However, one should remember that a typical CT of the lumbar spine will expose the patient to an effective dose of 5–10 millisieverts (mSv), which would be equivalent to 2.5–5 years of natural background radiation; those who travel on 7-hour flights (0.05 /uni00A0 mSv per 7-hour flight) would need to make 100–200 journeys in their lifetime to be exposed to the same effective dose.

---

Revision #1

Created 2025-12-31 15:15:08 UTC by Omar Ayman

Updated 2025-12-31 15:15:08 UTC by Omar Ayman