

# Cytology specimen

## Cytology specimen

Samples for cytology can be smeared immediately onto glass slides, fixed (usually in alcohol) or air dried and stained immediately or later. The process usually produces several slides, some of which are stained with a Papanicolaou (Pap) stain and some with another method such as May-Grünwald-Giemsa (MGG), H&E or Romanowsky ( Figure 11.9 ). Liquid-based thin-layer technology is now replacing older methods. For liquid-based cytology , the sampling device is usually washed in a liquid medium and the material obtained is then processed in the laboratory using purpose-built equipment. Cytology specimen

Samples for cytology can be smeared immediately onto glass slides, fixed (usually in alcohol) or air dried and stained immediately or later. The process usually produces several slides, some of which are stained with a Papanicolaou (Pap) stain and some with another method such as May-Grünwald-Giemsa (MGG), H&E or Romanowsky ( Figure 11.9 ). Liquid-based thin-layer technology is now replacing older methods. For liquid-based cytology , the sampling device is usually washed in a liquid medium and the material obtained is then processed in the laboratory using purpose-built equipment. Cytology specimen

Samples for cytology can be smeared immediately onto glass slides, fixed (usually in alcohol) or air dried and stained immediately or later. The process usually produces several slides, some of which are stained with a Papanicolaou (Pap) stain and some with another method such as May-Grünwald-Giemsa (MGG), H&E or Romanowsky ( Figure 11.9 ). Liquid-based thin-layer technology is now replacing older methods. For liquid-based cytology , the sampling device is usually washed in a liquid medium and the material obtained is then processed in the laboratory using purpose-built equipment.

---

Revision #1

Created 2025-12-31 15:08:18 UTC by Omar Ayman

Updated 2025-12-31 15:08:18 UTC by Omar Ayman