

# Danger to peripheral circulation

## Danger to peripheral circulation

In full-thickness burns, the collagen fibres are coagulated. The normal elasticity of the skin is lost. A circumferential full-thickness burn to a limb acts as a tourniquet as the limb swells. If untreated, this will progress to limb-threatening ischaemia. Summary box 46.7 Other complications of burns

Infection from the burn site, lungs, gut, lines and catheters  
Malabsorption from the gut  
Circumferential burns may compromise circulation to a limb

## Danger to peripheral circulation

In full-thickness burns, the collagen fibres are coagulated. The normal elasticity of the skin is lost. A circumferential full-thickness burn to a limb acts as a tourniquet as the limb swells. If untreated, this will progress to limb-threatening ischaemia. Summary box 46.7 Other complications of burns

Infection from the burn site, lungs, gut, lines and catheters  
Malabsorption from the gut  
Circumferential burns may compromise circulation to a limb

## Danger to peripheral circulation

In full-thickness burns, the collagen fibres are coagulated. The normal elasticity of the skin is lost. A circumferential full-thickness burn to a limb acts as a tourniquet as the limb swells. If untreated, this will progress to limb-threatening ischaemia. Summary box 46.7 Other complications of burns

Infection from the burn site, lungs, gut, lines and catheters  
Malabsorption from the gut  
Circumferential burns may compromise circulation to a limb

---

Revision #1

Created 2025-12-31 15:17:24 UTC by Omar Ayman

Updated 2025-12-31 15:17:24 UTC by Omar Ayman