

DECISION MAKING WITHIN THE DEPLOYED ENVIRONMENT

DECISION MAKING WITHIN THE DEPLOYED ENVIRONMENT

The damage control approach is vital in a proportion of war injuries, but thought must be given to available resources. Within a well-established role 3 unit, both DCS and definitive procedures are likely to be possible. Blood and blood products are also likely to be available and restocked regularly. In this mode of care. In contrast, decision making becomes more crucial within the austere environment. Theatre space, intensive care beds, - equipment and blood products may be limited. Long evacuation times may contraindicate permissive hypotension. Limited resources, such as blood, may need to be divided among casualties. Within a role 1 or role 2 facility, immediate evacuation may be undertaken in preference to more thorough stabilisation. Decision making of this nature is complex and difficult and requires practice in the context of simulation, exercises and courses. - DECISION MAKING WITHIN THE DEPLOYED ENVIRONMENT

The damage control approach is vital in a proportion of war injuries, but thought must be given to available resources. Within a well-established role 3 unit, both DCS and definitive procedures are likely to be possible. Blood and blood products are also likely to be available and restocked regularly. In this mode of care. In contrast, decision making becomes more crucial within the austere environment. Theatre space, intensive care beds, - equipment and blood products may be limited. Long evacuation times may contraindicate permissive hypotension. Limited resources, such as blood, may need to be divided among casualties. Within a role 1 or role 2 facility, immediate evacuation may be undertaken in preference to more thorough stabilisation. Decision making of this nature is complex and difficult and requires practice in the context of simulation, exercises and courses. - DECISION MAKING WITHIN THE DEPLOYED ENVIRONMENT

The damage control approach is vital in a proportion of war injuries, but thought must be given to available resources. Within a well-established role 3 unit, both DCS and definitive procedures are likely to be possible. Blood and blood products are also likely to be available and restocked regularly. In this mode of care. In contrast, decision making becomes more crucial within the austere environment. Theatre space, intensive care beds, - equipment and blood products may be limited. Long evacuation times may contraindicate permissive hypotension. Limited resources, such as blood, may need to be divided among casualties. Within a role 1 or role 2 facility, immediate evacuation may be undertaken in preference to more thorough stabilisation. Decision making of this nature is complex and difficult and requires practice in the context of simulation, exercises and courses. -

Revision #1

Created 2025-12-31 15:14:07 UTC by Omar Ayman

Updated 2025-12-31 15:14:07 UTC by Omar Ayman