

# Dermis

## Dermis

The dermis constitutes 95% of the skin and is structurally divided into a superficial papillary layer, which is composed of delicate collagen and elastin fibres in ground substance, into which a capillary and lymphatic network ramifies, and a deeper reticular layer, which is composed of coarse branching collagen, layered parallel to the skin surface ( Figure 45.1 ). - The epidermis and dermis meet at the dermoepidermal junction in a three-dimensional wave-like arrangement in - which epidermal rete ridges project down, inter digitating with the upward-pointing, dermal papillae containing vascular and lymphatic plexi. m - The skin also contains specialised cells such as Langerhans cells, whose role is to engulf antigens and present them to T cells. Merkel cells, and Meissner's and Pacinian corpuscles have roles in mechanosensation. - Dermis

The dermis constitutes 95% of the skin and is structurally divided into a superficial papillary layer, which is composed of delicate collagen and elastin fibres in ground substance, into which a capillary and lymphatic network ramifies, and a deeper reticular layer, which is composed of coarse branching collagen, layered parallel to the skin surface ( Figure 45.1 ). - The epidermis and dermis meet at the dermoepidermal junction in a three-dimensional wave-like arrangement in - which epidermal rete ridges project down, inter digitating with the upward-pointing, dermal papillae containing vascular and lymphatic plexi. m - The skin also contains specialised cells such as Langerhans cells, whose role is to engulf antigens and present them to T cells. Merkel cells, and Meissner's and Pacinian corpuscles have roles in mechanosensation. - Dermis

The dermis constitutes 95% of the skin and is structurally divided into a superficial papillary layer, which is composed of delicate collagen and elastin fibres in ground substance, into which a capillary and lymphatic network ramifies, and a deeper reticular layer, which is composed of coarse branching collagen, layered parallel to the skin surface ( Figure 45.1 ). - The epidermis and dermis meet at the dermoepidermal junction in a three-dimensional wave-like arrangement in - which epidermal rete ridges project down, inter digitating with the upward-pointing, dermal papillae containing vascular and lymphatic plexi. m - The skin also contains specialised cells such as Langerhans cells, whose role is to engulf antigens and present them to T cells. Merkel cells, and Meissner's and Pacinian corpuscles have roles in mechanosensation. -

---

Revision #1

Created 2025-12-31 15:17:04 UTC by Omar Ayman

Updated 2025-12-31 15:17:04 UTC by Omar Ayman