

Discogenic low back pain

Discogenic low back pain

Discogenic low back pain has been defined as a continuum of diagnostic categories (internal disc disruption, degenerative disc disease, segmental instability) reflecting various stages of degenerative pathology affecting the intervertebral disc. Not all degenerate discs are painful. Patients typically present with chronic relapsing episodes of low back pain between the ages of 40 and 60 years. A recent study has compared rehabilitation with spinal fusion for discogenic pain. Both groups reported reductions in disability, with the authors strongly recommending a course of rehabilitation before surgical intervention. For those who fail to improve with conservative measures, provocative lumbar discography (Figure 37.1) may help to identify the source of pain, and surgery in the form of a lumbar spinal fusion - (Figure 37.3) or lumbar disc replacement (Figure 37.4) may be considered. J E Buck, surgeon, Brook General Hospital, London, UK, described the direct repair of the defect in spondylolisthesis in 1970.

(b) (c) Figure 37.3 Anterior lumbar interbody fusion. (a) The PEEK (poly-ethyl-ethyl-ketone) cage has been packed with bone graft prior to insertion; (b, c) show the anteroposterior and lateral postoperative radiographs, respectively. (b) Figure 37.4 Lumbar total disc replacement. (a)

Anteroposterior radio

graph with 30° of cranial inclination. (b) Lateral radiograph with the implant appropriately positioned.

Discogenic low back pain

Discogenic low back pain has been defined as a continuum of diagnostic categories (internal disc disruption, degenerative disc disease, segmental instability) reflecting various stages of degenerative pathology affecting the intervertebral disc. Not all degenerate discs are painful. Patients typically present with chronic relapsing episodes of low back pain between the ages of 40 and 60 years. A recent study has compared rehabilitation with spinal fusion for discogenic pain. Both groups reported reductions in disability, with the authors strongly recommending a course of rehabilitation before surgical intervention. For those who fail to improve with conservative measures, provocative lumbar discography (Figure 37.1) may help to identify the source of pain, and surgery in the form of a lumbar spinal fusion - (Figure 37.3) or lumbar disc replacement (Figure 37.4) may be considered. J E Buck, surgeon, Brook General Hospital, London, UK, described the direct repair of the defect in spondylolisthesis in 1970.

(b) (c) Figure 37.3 Anterior lumbar interbody fusion. (a) The PEEK (poly-ethyl-ethyl-ketone) cage has been packed with bone graft prior to insertion; (b, c) show the anteroposterior and lateral postoperative radiographs, respectively. (b) Figure 37.4

Lumbar total disc replacement. (a) Anteroposterior radio

graph with 30° of cranial inclination. (b) Lateral radiograph with the implant appropriately positioned.

Discogenic low back pain

Discogenic low back pain has been defined as a continuum of diagnostic categories (internal disc disruption, degenerative disc disease, segmental instability) reflecting various stages of degenerative pathology affecting the intervertebral disc. Not all degenerate discs are painful. Patients typically present with chronic relapsing episodes of low back pain between the ages of 40 and 60 years. A recent study has compared rehabilitation with spinal fusion for discogenic pain. Both groups reported reductions in disability, with the authors strongly recommending a course of rehabilitation before surgical intervention. For those who fail to improve with conservative measures, provocative lumbar discography (Figure 37.1) may help to identify the source of pain, and surgery in the form of a lumbar spinal fusion - (Figure 37.3) or lumbar disc replacement (Figure 37.4) may be considered. J E Buck, surgeon, Brook General Hospital, London, UK, described the direct repair of the defect in spondylolisthesis in 1970.

(b) (c) Figure 37.3 Anterior lumbar interbody fusion. (a) The PEEK (poly-ethyl-ethyl-ketone) cage has been packed with bone graft prior to insertion; (b, c) show the anteroposterior and lateral postoperative radiographs,

respectively. (b) Figure 37.4 Lumbar total disc replacement. (a) Anteroposterior radio

graph with 30° of cranial inclination. (b) Lateral radiograph with the implant appropriately positioned.

Revision #1

Created 2025-12-31 15:15:10 UTC by Omar Ayman

Updated 2025-12-31 15:15:10 UTC by Omar Ayman