

Follicular carcinoma

Follicular carcinoma

Follicular carcinoma can normally only be differentiated from follicular adenoma by the architecture on histology . For this reason, follicular lesions on FNA are unable to be diagnosed as malignant in the absence of clinical features such as metastases - (Figure 55.24). Multiple foci of follicular carcinoma are seldom seen and lymph node involvement is much less common than in PTC. Blood-borne metastases are more common and the - eventual mortality rate, although still low , is twice that of PTC (Figure 55.25). Hürthle cell tumours are a rare variant of follicular neoplasm in which oxyphil (Hürthle, Askanazy) cells predominate histologically . Hürthle cell cancers are associated with a poor prognosis. - Follicular carcinoma

Follicular carcinoma can normally only be differentiated from follicular adenoma by the architecture on histology . For this reason, follicular lesions on FNA are unable to be diagnosed as malignant in the absence of clinical features such as metastases - (Figure 55.24). Multiple foci of follicular carcinoma are seldom seen and lymph node involvement is much less common than in PTC. Blood-borne metastases are more common and the - eventual mortality rate, although still low , is twice that of PTC (Figure 55.25). Hürthle cell tumours are a rare variant of follicular neoplasm in which oxyphil (Hürthle, Askanazy) cells predominate histologically . Hürthle cell cancers are associated with a poor prognosis. -

Revision #1

Created 2025-12-31 15:20:48 UTC by Omar Ayman

Updated 2025-12-31 15:20:48 UTC by Omar Ayman