

FURTHER READING

FURTHER READING

Busuttil R, Klimm G (eds). Transplantation of the liver , 3rd edn. Philadelphia, PA: Elsevier-Saunders, 2015. Clavien P , Trotter J (eds). Medical care of the liver transplant patient, edn. Oxford: Wiley-Blackwell, 2012. Croome K, Muiesan P , Taner B (eds). Donation after circulatory death (DCD) liver transplantation: a practical guide . Springer, 2020. Fan ST (ed.). Living donor liver transplantation, 2nd edn. Singapore: World Scientific, 2011. Hadzic N, Baumann U, McLin V . Pediatric liver transplantation: a clinical guide . Philadelphia, PA: Elsevier, 2021. Hricik D (ed.). Primer on transplantation, 3rd edn. Oxford: Wiley- Blackwell, 2011. Garden OJ, Parks RW , Wigmore SJ (eds). Principles and practice of surgery, 8th edn. Elsevier, 2022. Oniscu G, Forsythe J, Fung J (eds) . Abdominal organ retrieval and transplantation bench surgery . Oxford: Wiley-Blackwell, 2013. Sutcliffe R, Antoniadou C, Deshpande R et al . Liver and pancreatobiliary surgery: with liver transplantation. Oxford: Oxford University Press, 2009. Definitions /uni25CF /uni25CF /uni25CF - /uni25CF /uni25CF /uni25CF /uni25CF - /uni25CF /uni25CF

Acute liver failure: complex multisystem illness that evolves quickly after a catastrophic insult to the liver, leading to coagulopathy and encephalopathy; based on onset of encephalopathy – hyperacute (within 7 days), acute (8–28 days), subacute (>28 days) Allograft: an organ or tissue transplanted from one individual to another Chronic liver disease: progressive deterioration of liver functions for more than 6 months, with in /f_l ammation, destruction and regeneration of liver parenchyma, leading to /f_i brosis and cirrhosis Heterotopic graft: a graft placed in a site different from that where the organ is normally located HLA: human leukocyte antigen, the main trigger to graft rejection MELD score: model for end-stage liver disease score; predicts prognosis in patients with chronic liver disease and is used to allocate liver for transplant NAFLD: non-alcoholic fatty liver disease; associated with metabolic syndrome (obesity, hyperlipidaemia and diabetes mellitus). Some patients develop non-alcoholic steatohepatitis (NASH), which leads to /f_i brosis and cirrhosis Orthotopic graft: a graft placed in its normal anatomical site UKELD score: United Kingdom model for end-stage liver disease score; predicts survival of patients listed for LT in the UK

Revision #1

Created 2025-12-31 15:31:49 UTC by Omar Ayman

Updated 2025-12-31 15:31:49 UTC by Omar Ayman