

FURTHER READING

FURTHER READING

Bainbridge D, Martin J, Arango M, Cheng D; for the Evidence-based Peri-operative Clinical Outcomes Research (EPiCOR) Group. Perioperative and anaesthetic-related mortality in developed and developing countries: a systematic review and meta-analysis. *Lancet* 2012; 380 (9847): 1075–81.

Barker P , Creasey PE, Dhatariya K et al . Peri-operative management of the surgical patient with diabetes. *Anaesthesia* 2015; 70 (12): 1427–40.

Boyd O, Jackson N. How is risk defined in high-risk surgical patient management? *Crit Care* 2005; 9 : 390–6.

Chan MTV , Wang CY , Edwin S et al . Association of unrecognized obstructive sleep apnoea with postoperative cardiovascular events in patients undergoing major non cardiac surgery . *JAMA* 2019; 321 (18): 1788–98.

Department of Health. Mental Capacity Act (MCA). England and Wales . London: HMSO, 2005.

Duminda N, Wijesundera W , Beattie S et al . Integration of the Duke Activity Status Index into preoperative risk evaluation: a multicentre prospective cohort study . *Br J Anaesth* 2020; 124 (3): 261–70.

Fleisher L, Fleischmann K, Auerback A et al . 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014; 64 (22): e77–e137. from <https://www.gmc-uk.org/ethical-guidance/ethical-guidance-for-doctors/decision-making-and-consent>.

Gri ffi ths R, Beech F , Brown A et al . Peri-operative care of the elderly 2014. *Anaesthesia*. 2014; 69 : 81–98.

Gri ffi ths R, Babu S, Dixon S et al . Guideline for the management of hip fractures 2020. *Anaesthesia* 2021; 76 : 225–37.

Hartle A, McCormack T , Carlisle J et al . The measurement of adult blood pressure and management of hypertension before elective surgery . *Anaesthesia* 2016; 71 (3): 326–37.

Jørgensen ME, Torp-Pedersen C, Gislason GH et al . Time elapsed after ischemic stroke and risk of adverse cardiovascular events and mortality following elective noncardiac surgery . *JAMA* 312 (3): 269–77.

Kristenson SD, Knuuti J, Saraste A et al . 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *Eur Heart J* 2014; 35 : 2383–431.

Lee TH, Marcantonio ER, Mangione CM et al . Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery . *Circulation* 1999; 100 : 1043–9.

Minto G, Biccard B. Assessment of the high-risk perioperative patient. *BJA Educ* 2014; 14 (1): 12–17.

Munoz M, Acheson G, Auerbach M et al . International consensus statement on the perioperative management of anaemia and iron deficiency . *Anaesthesia* 2017; 72 (2): 233–47.

Murray P , Whiting P , Hutchinson S et al . Preoperative shuttle walking testing and outcome after oesophagogastrrectomy . *Br J Anaesth* 2007; 99 (6): 809–11.

National Confidential Enquiry into Patient Outcome and Death (NCEPOD). An age-old problem: a review of the care received by elderly patients undergoing surgery, 2010. Available from https://www.ncepod.org.uk/2010report3/downloads/EESE_fullReport.pdf.

National Confidential Enquiry into Patient Outcome and Death (NCEPOD). Knowing the risk. A review of the perioperative care of surgical patients , 2011. Available from <https://www.ncepod.org>.

uk/2011report2/downloads/POC_fullreport.pdf. National Emergency Laparotomy Audit (NELA) Project Team. patient report of the National Emergency Laparotomy Audit , 2020. Available from <https://www.nela.org.uk/reports>. tests for elective surgery . NICE Guideline 45. London: NICE, 2016. Available from <https://www.nice.org.uk/guidance/ng45>. National Institute for Health and Care Excellence. Decision-making and mental capacity . NICE Guideline 108. London: NICE, 2018. Available from <https://www.nice.org.uk/guidance/ng108>. National Institute for Health and Care Excellence. Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism . NICE Guideline 89. London: NICE, 2019. Available from <https://www.nice.org.uk/guidance/ng89>. National Institute for Health and Care Excellence. Perioperative care in adults 2014; . NICE Guideline 180. London: NICE, 2020. Available from <https://www.nice.org.uk/guidance/ng180>. Noordzij PG, Poldermans D, Schouten O et al . Postoperative mortality in The Netherlands: a population based analysis of surgery-specific risk in adults. *Anesthesiology* 2010; 112 (5): 1105–15. Practice guidelines for the perioperative management of patients with obstructive sleep apnea: a report by the American Society of Anesthesiologists Task Force on Perioperative Management of Patients with Obstructive Sleep Apnea. *Anesthesiology* 2006; 104 : 1081–93. Royal College of Surgeons. Caring for patients who refuse blood: a guide to good practice for the surgical management of Jehovah's Witnesses and other patients who decline transfusion , 2016. Available from <https://www.rcseng.ac.uk/-/media/files/rcs/library-and-publications/non-journal-publications/caring-for-patients-who-refuse-blood-a-guide-to-good-practice.pdf>. Royal College of Surgeons. Consent: supported decision making: a guide to good practice , 2018. Available from <https://www.rcseng.ac.uk/standards-and-research/standards-and-guidance/good-practice-guides/consent/>. Shoemaker WC, Appel PL, Kram HB et al . Hemodynamic and oxygen transport responses in survivors and non survivors of high risk surgery . *Crit Care Med* 1993; 21 (7): 977–90. Woodcock T , Barker P , Daniel S et al . Guidelines for the management of glucocorticoids during the perioperative period for patients with adrenal insufficiency . *Anaesthesia* 2020; 75 (5): 654–63. Sixth FURTHER READING

Bainbridge D, Martin J, Arango M, Cheng D; for the Evidence-based Peri-operative Clinical Outcomes Research (EPiCOR) Group. Perioperative and anaesthetic-related mortality in developed and developing countries: a systematic review and meta-analysis. *Lancet* 2012; 380 (9847): 1075–81. Barker P , Creasey PE, Dhatariya K et al . Peri-operative management of the surgical patient with diabetes. *Anaesthesia* 2015; 70 (12): 1427–40. Boyd O, Jackson N. How is risk defined in high-risk surgical patient management? *Crit Care* 2005; 9 : 390–6. Chan MTV , Wang CY , Edwin S et al . Association of unrecognized obstructive sleep apnoea with postoperative cardiovascular events in patients undergoing major non cardiac surgery . *JAMA* 2019; 321 (18): 1788–98. Department of Health. Mental Capacity Act (MCA). England and Wales . London: HMSO, 2005. Duminda N, Wijesundera W , Beattie S et al . Integration of the Duke Activity Status Index into preoperative risk evaluation: a multicentre prospective cohort study . *Br J Anaesth* 2020; 124 (3): 261–70. Fleisher L, Fleischmann K, Auerback A et al . 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014; 64 (22): e77–e137. from <https://www.gmc-uk.org/ethical-guidance/ethical-guidance-for-doctors/decision-making-and-consent>. Griffiths R, Beech F , Brown A et al . Perioperative care of the elderly 2014. *Anaesthesia*. 2014; 69 : 81–98. Griffiths R, Babu S, Dixon S et al . Guideline for the management of hip fractures 2020. *Anaesthesia* 2021; 76 : 225–37. Hartle A, McCormack T , Carlisle J et al . The measurement of adult blood pressure and management of hypertension before elective surgery . *Anaesthesia* 2016; 71 (3): 326–37. Jørgensen ME, Torp-

Pedersen C, Gislason GH et al . Time elapsed after ischemic stroke and risk of adverse cardiovascular events and mortality following elective noncardiac surgery . JAMA 312 (3): 269–77.

Kristenson SD, Knuuti J, Saraste A et al . 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). Eur Heart J 2014; 35 : 2383–431.

Lee TH, Marcantonio ER, Mangione CM et al . Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery . Circulation 1999; 100 : 1043–9.

Minto G, Biccard B. Assessment of the high-risk perioperative patient. BJA Educ 2014; 14 (1): 12–17.

Munoz M, Acheson G, Auerbach M et al . International consensus statement on the perioperative management of anaemia and iron deficiency . Anaesthesia 2017; 72 (2): 233–47.

Murray P , Whiting P , Hutchinson S et al . Preoperative shuttle walking testing and outcome after oesophagogastrrectomy . Br J Anaesth 2007; 99 (6): 809–11.

National Confidential Enquiry into Patient Outcome and Death (NCEPOD). An age-old problem: a review of the care received by elderly patients undergoing surgery, 2010. Available from https://www.ncepod.org.uk/2010report3/downloads/EESE_fullReport.pdf.

National Confidential Enquiry into Patient Outcome and Death (NCEPOD). Knowing the risk. A review of the perioperative care of surgical patients , 2011. Available from https://www.ncepod.org.uk/2011report2/downloads/POC_fullreport.pdf.

National Emergency Laparotomy Audit (NELA) Project Team. patient report of the National Emergency Laparotomy Audit , 2020. Available from <https://www.nela.org.uk/reports>.

tests for elective surgery . NICE Guideline 45. London: NICE, 2016. Available from <https://www.nice.org.uk/guidance/ng45>.

National Institute for Health and Care Excellence. Decision-making and mental capacity . NICE Guideline 108. London: NICE, 2018. Available from <https://www.nice.org.uk/guidance/ng108>.

National Institute for Health and Care Excellence. Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism . NICE Guideline 89. London: NICE, 2019. Available from <https://www.nice.org.uk/guidance/ng89>.

National Institute for Health and Care Excellence. Perioperative care in adults 2014; . NICE Guideline 180. London: NICE, 2020. Available from <https://www.nice.org.uk/guidance/ng180>.

Noordzij PG, Poldermans D, Schouten O et al . Postoperative mortality in The Netherlands: a population based analysis of surgery-specific risk in adults. Anesthesiology 2010; 112 (5): 1105–15.

Practice guidelines for the perioperative management of patients with obstructive sleep apnea: a report by the American Society of Anesthesiologists Task Force on Perioperative Management of Patients with Obstructive Sleep Apnea. Anesthesiology 2006; 104 : 1081–93.

Royal College of Surgeons. Caring for patients who refuse blood: a guide to good practice for the surgical management of Jehovah’s Witnesses and other patients who decline transfusion , 2016. Available from <https://www.rcseng.ac.uk/-/media/files/rcs/library-and-publications/non-journal-publications/caring-for-patients-who-refuse-blood--a-guide-to-good-practice.pdf>.

Royal College of Surgeons. Consent: supported decision making: a guide to good practice , 2018. Available from <https://www.rcseng.ac.uk/standards-and-research/standards-and-guidance/good-practice-guides/consent/>.

Shoemaker WC, Appel PL, Kram HB et al . Hemodynamic and oxygen transport responses in survivors and non survivors of high risk surgery . Crit Care Med 1993; 21 (7): 977–90.

Woodcock T , Barker P , Daniel S et al . Guidelines for the management of glucocorticoids during the perioperative period for patients with adrenal insufficiency . Anaesthesia 2020; 75 (5): 654–63.

