

FURTHER READING

FURTHER READING

Al-Batran S-E, Homann N, Schmalenberg H et al . Perioperative chemotherapy with docetaxel, oxaliplatin, and fluorouracil/leucovorin (FLOT) versus epirubicin, cisplatin, and fluorouracil or capecitabine (ECF/ECX) for resectable gastric or gastroesophageal junction (GEJ) adenocarcinoma (FLOT4-AIO): a multicenter, randomized phase 3 trial. *J Clin Oncol* 2017; 35 (15 suppl): 4004. Cristescu R, Lee J, Nebozhyn M et al . Molecular analysis of gastric cancer identifies subtypes associated with distinct clinical outcomes. *Nat Med* 2015; 21 (5): 449-56. Lee SS, Chung HY , Kwon OK, Yu W . Long-term quality of life after distal subtotal and total gastrectomy: symptom- and behaviour - oriented consequences. *Ann Surg* 2016; 263 (4): 738-44. Smyth EC, Nilsson M, Grabsch HI et al . Gastric cancer. *Lancet* 2020; 396 (10251): 635-48. Wilson MS, Blencowe NS, Boyle C et al. ; AUGIS. A modified Delphi process to establish future research priorities in malignant oesophageal - agastric surgery . *Surgeon* 2020; 18 (6): 321-6.

•

Revision #1

Created 2025-12-31 15:24:55 UTC by Omar Ayman

Updated 2025-12-31 15:24:55 UTC by Omar Ayman