

# Gastrointestinal haemorrhage

## Gastrointestinal haemorrhage

The aetiology of acute gastrointestinal (GI) haemorrhage varies between the upper GIT (common causes including peptic ulcer disease, varices and Mallory-Weiss tears) and the lower GIT (common causes including angiodysplasia, diverticular haemorrhage and neoplastic lesions). While endoscopy is a useful first-line investigation for both, in refractory or occult GI haemorrhage radiology can also contribute to diagnosis and management. Nuclear medicine scans using radioisotope-labelled red blood cells are useful when bleeding is intermittent, but for patients suspected of active bleeding the best investigation is a CT mesenteric angiogram. Non-contrast scans to look for bright blood in the bowel lumen should be supplemented with scans in the arterial phase to assess for a blush due to active extravasation and the portal venous phase to optimise detection of wall thickening and masses and to look for sites of venous bleeding. If non-invasive imaging is effective, catheter angiography can be used to embolise a bleeding point.

## Gastrointestinal haemorrhage

The aetiology of acute gastrointestinal (GI) haemorrhage varies between the upper GIT (common causes including peptic ulcer disease, varices and Mallory-Weiss tears) and the lower GIT (common causes including angiodysplasia, diverticular haemorrhage and neoplastic lesions). While endoscopy is a useful first-line investigation for both, in refractory or occult GI haemorrhage radiology can also contribute to diagnosis and management. Nuclear medicine scans using radioisotope-labelled red blood cells are useful when bleeding is intermittent, but for patients suspected of active bleeding the best investigation is a CT mesenteric angiogram. Non-contrast scans to look for bright blood in the bowel lumen should be supplemented with scans in the arterial phase to assess for a blush due to active extravasation and the portal venous phase to optimise detection of wall thickening and masses and to look for sites of venous bleeding. If non-invasive imaging is effective, catheter angiography can be used to embolise a bleeding point.

## Gastrointestinal haemorrhage

The aetiology of acute gastrointestinal (GI) haemorrhage varies between the upper GIT (common causes including peptic ulcer disease, varices and Mallory-Weiss tears) and the lower GIT (common causes including angiodysplasia, diverticular haemorrhage and neoplastic lesions). While endoscopy is a useful first-line investigation for both, in refractory or occult GI haemorrhage radiology can also contribute to diagnosis and management. Nuclear medicine scans using radioisotope-labelled red blood cells are useful when bleeding is intermittent, but for patients suspected of active bleeding the best investigation is a CT mesenteric angiogram. Non-contrast scans to look for bright blood in the bowel lumen should be supplemented with scans in the arterial phase to assess for a blush due to active extravasation and the portal venous phase to optimise detection of wall thickening and masses and to look for sites of venous bleeding. If non-invasive

imaging is effective, catheter angiography can be used to embolise a bleeding point.

---

Revision #1

Created 2025-12-31 15:28:46 UTC by Omar Ayman

Updated 2025-12-31 15:28:46 UTC by Omar Ayman