

# HAEMORRHAGE RESUSCITATION

## HAEMORRHAGE RESUSCITATION

The conduct and goals of resuscitation change depending on whether the patient is actively bleeding. In this case, the resuscitation focuses on achieving rapid haemostasis and maintaining the ability of the blood to clot. This paradigm is called damage control resuscitation (see Damage control resuscitation If the patient is not actively bleeding, has stopped bleeding or the cause of shock is not haemorrhage, then resuscitation is directed at correcting the shock state and restoring perfusion to end organs. HAEMORRHAGE RESUSCITATION

The conduct and goals of resuscitation change depending on whether the patient is actively bleeding. In this case, the resuscitation focuses on achieving rapid haemostasis and maintaining the ability of the blood to clot. This paradigm is called damage control resuscitation (see Damage control resuscitation If the patient is not actively bleeding, has stopped bleeding or the cause of shock is not haemorrhage, then resuscitation is directed at correcting the shock state and restoring perfusion to end organs. HAEMORRHAGE RESUSCITATION

The conduct and goals of resuscitation change depending on whether the patient is actively bleeding. In this case, the resuscitation focuses on achieving rapid haemostasis and maintaining the ability of the blood to clot. This paradigm is called damage control resuscitation (see Damage control resuscitation If the patient is not actively bleeding, has stopped bleeding or the cause of shock is not haemorrhage, then resuscitation is directed at correcting the shock state and restoring perfusion to end organs.

---

Revision #1

Created 2025-12-31 15:10:15 UTC by Omar Ayman

Updated 2025-12-31 15:10:15 UTC by Omar Ayman