

Immobility

Immobility

Immobility has long been recognised as a potent stimulus for inducing muscle wasting. Inactivity impairs the normal meal-derived amino acid stimulation of protein synthesis in skeletal muscle. Avoidance of unnecessary bed rest and active early mobilisation are essential measures to avoid muscle wasting as a consequence of immobility . Pre-habilitation programmes provide a better starting point before surgery . Immobility

Immobility has long been recognised as a potent stimulus for inducing muscle wasting. Inactivity impairs the normal meal-derived amino acid stimulation of protein synthesis in skeletal muscle. Avoidance of unnecessary bed rest and active early mobilisation are essential measures to avoid muscle wasting as a consequence of immobility . Pre-habilitation programmes provide a better starting point before surgery . Immobility

Immobility has long been recognised as a potent stimulus for inducing muscle wasting. Inactivity impairs the normal meal-derived amino acid stimulation of protein synthesis in skeletal muscle. Avoidance of unnecessary bed rest and active early mobilisation are essential measures to avoid muscle wasting as a consequence of immobility . Pre-habilitation programmes provide a better starting point before surgery .

Revision #1

Created 2025-12-31 15:07:43 UTC by Omar Ayman

Updated 2025-12-31 15:07:43 UTC by Omar Ayman