

Knock knees and bowlegs

Knock knees and bowlegs

All children start life with bowlegs, often accompanied by internal tibial torsion. By the age of 2-3 years they have developed knock knees, which regress towards the normal adult tibiofemoral angle of 7° valgus by age 7 (Figure 44.3). The intercondylar or intermalleolar distance is often used to quantify the deformity but radiographs are needed when the deformity is severe, asymmetrical or symptomatic. The most common pathological causes are previous trauma, rickets or a skeletal dysplasia. Knock knees and bowlegs

All children start life with bowlegs, often accompanied by internal tibial torsion. By the age of 2-3 years they have developed knock knees, which regress towards the normal adult tibiofemoral angle of 7° valgus by age 7 (Figure 44.3). The intercondylar or intermalleolar distance is often used to quantify the deformity but radiographs are needed when the deformity is severe, asymmetrical or symptomatic. The most common pathological causes are previous trauma, rickets or a skeletal dysplasia. Knock knees and bowlegs

All children start life with bowlegs, often accompanied by internal tibial torsion. By the age of 2-3 years they have developed knock knees, which regress towards the normal adult tibiofemoral angle of 7° valgus by age 7 (Figure 44.3). The intercondylar or intermalleolar distance is often used to quantify the deformity but radiographs are needed when the deformity is severe, asymmetrical or symptomatic. The most common pathological causes are previous trauma, rickets or a skeletal dysplasia.

Revision #1

Created 2025-12-31 15:16:47 UTC by Omar Ayman

Updated 2025-12-31 15:16:47 UTC by Omar Ayman