

# Lacrimal system

## Lacrimal system

The almond-shaped lacrimal gland lies under the upper outer orbital rim and opens into the upper conjunctival fornix through 10–15 ducts. Tears are swept across the globe by the lids and evaporate or pass into the upper and lower lid puncta, and then into the canaliculi to join the common canaliculus, which passes into the lacrimal sac under the medial canthal tendon. The sac is drained by the nasolacrimal duct into the nose, opening in the inferior meatus under the inferior turbinate. Lacrimal system

The almond-shaped lacrimal gland lies under the upper outer orbital rim and opens into the upper conjunctival fornix through 10–15 ducts. Tears are swept across the globe by the lids and evaporate or pass into the upper and lower lid puncta, and then into the canaliculi to join the common canaliculus, which passes into the lacrimal sac under the medial canthal tendon. The sac is drained by the nasolacrimal duct into the nose, opening in the inferior meatus under the inferior turbinate. Lacrimal system

The almond-shaped lacrimal gland lies under the upper outer orbital rim and opens into the upper conjunctival fornix through 10–15 ducts. Tears are swept across the globe by the lids and evaporate or pass into the upper and lower lid puncta, and then into the canaliculi to join the common canaliculus, which passes into the lacrimal sac under the medial canthal tendon. The sac is drained by the nasolacrimal duct into the nose, opening in the inferior meatus under the inferior turbinate.

---

Revision #1

Created 2025-12-31 15:18:13 UTC by Omar Ayman

Updated 2025-12-31 15:18:13 UTC by Omar Ayman