

Limb salvage

Limb salvage

The Mangled Extremity Severity Score (MESS) and its modifications are useful in deciding about limb salvage. Extensive tissue loss, neurovascular damage and loss of long fragments of bone are traditionally indications for amputation. Currently, wounds of any dimension can be covered with microvascular flaps and distraction osteogenesis and vascularised bone can be used to restore bony continuity. If performed in time, vascular repairs can salvage most acutely ischaemic limbs. Because of these developments the indications for amputation in trauma have undergone a paradigm shift and the majority of patients who reach a tertiary-care facility within 24 hours are candidates for limb salvage (Figure 33.13). This assumes that debridement and, if required, vascular repairs have been performed in a field medical facility. A limb is unlikely to survive if the vascular repair of major limb vessels has been delayed for more than 4-6 hours.

(c) Figure 33.13 (a-d) Badly traumatised lower limb. Reconstruction has been performed using a microvascular rectus abdominis flap covered with a skin graft.

Limb salvage

The Mangled Extremity Severity Score (MESS) and its modifications are useful in deciding about limb salvage. Extensive tissue loss, neurovascular damage and loss of long fragments of bone are traditionally indications for amputation. Currently, wounds of any dimension can be covered with microvascular flaps and distraction osteogenesis and vascularised bone can be used to restore bony continuity. If performed in time, vascular repairs can salvage most acutely ischaemic limbs. Because of these developments the indications for amputation in trauma have undergone a paradigm shift and the majority of patients who reach a tertiary-care facility within 24 hours are candidates for limb salvage (Figure 33.13). This assumes that debridement and, if required, vascular repairs have been performed in a field medical facility. A limb is unlikely to survive if the vascular repair of major limb vessels has been delayed for more than 4-6 hours.

(c) Figure 33.13 (a-d) Badly traumatised lower limb. Reconstruction has been performed using a microvascular rectus abdominis flap covered with a skin graft.

Limb salvage

The Mangled Extremity Severity Score (MESS) and its modifications are useful in deciding about limb salvage. Extensive tissue loss, neurovascular damage and loss of long fragments of bone are traditionally indications for amputation. Currently, wounds of any dimension can be covered with microvascular flaps and distraction osteogenesis and vascularised bone can be used to restore bony continuity. If performed in time, vascular repairs can salvage most acutely ischaemic limbs.

Because of these developments the indications for amputation in trauma have undergone a paradigm shift and the majority of patients who reach a tertiary-care facility within 24 hours are candidates for limb salvage (Figure 33.13). This assumes that debridement and, if required, vascular repairs have been performed in a field medical facility . A limb is unlikely to survive if the vascular repair of major limb vessels has been delayed for more than 4–6 hours.

(c) Figure 33.13 (a–d) Badly traumatised lower limb. Reconstruction has been performed using a microvascular rectus abdominis flap covered with a skin graft.

Revision #1

Created 2025-12-31 15:13:57 UTC by Omar Ayman

Updated 2025-12-31 15:13:57 UTC by Omar Ayman