

Microbiology

Microbiology

Virulent Gram-positive organisms, particularly *Staphylococcus aureus*, are the most common cause of bone or joint infection in native tissue. However, once prosthetic material is implanted, a wide range of organisms can be involved. This includes - organisms with low virulence that are usually considered skin commensals, such as coagulase-negative staphylococci, α -haemolytic streptococci and *Cutibacterium acnes* (Table 43.1). e Microbiology

Virulent Gram-positive organisms, particularly *Staphylococcus aureus*, are the most common cause of bone or joint infection in native tissue. However, once prosthetic material is implanted, a wide range of organisms can be involved. This includes - organisms with low virulence that are usually considered skin commensals, such as coagulase-negative staphylococci, α -haemolytic streptococci and *Cutibacterium acnes* (Table 43.1). e Microbiology

Virulent Gram-positive organisms, particularly *Staphylococcus aureus*, are the most common cause of bone or joint infection in native tissue. However, once prosthetic material is implanted, a wide range of organisms can be involved. This includes - organisms with low virulence that are usually considered skin commensals, such as coagulase-negative staphylococci, α -haemolytic streptococci and *Cutibacterium acnes* (Table 43.1). e

Revision #1

Created 2025-12-31 15:16:32 UTC by Omar Ayman

Updated 2025-12-31 15:16:32 UTC by Omar Ayman