

Microscopic features of inflammation

Microscopic features of inflammation

Acute inflammation is characterised histologically by neutrophils (polymorphonuclear leukocytes), erosion or ulceration (Figure 11.18) and chronic inflammation by lymphocytes and plasma cells. Other inflammatory cells include eosinophils (Figure 11.19) , mast cells and histiocytes. Granulomas (collections of epithelioid histiocytes) (Figure 11.20a) raise the possibility of mycobacterial infection (Figure 11.20b), fungal 's disease or a reaction infection, parasites, sarcoidosis, Crohn to foreign material, among numerous other possible causes. s may reflect parasitic infection or Eosinophils in large number allergy . Interpretation depends heavily on the site and clinical setting.

Figure 11.18 An acute inflammatory process characterised by numerous neutrophils. Note the typical multilobated nuclei (arrows). Figure 11.19 Oesophageal mucosa infiltrated by numerous eosinophils with bright red cytoplasm, many of which are forming clusters. Eosinophils may reflect allergy, parasitic infection or a wide variety of other causes. In this example, the clinicopathological diagnosis was eosinophilic oesophagitis.

Microscopic features of inflammation

Acute inflammation is characterised histologically by neutrophils (polymorphonuclear leukocytes), erosion or ulceration (Figure 11.18) and chronic inflammation by lymphocytes and plasma cells. Other inflammatory cells include eosinophils (Figure 11.19) , mast cells and histiocytes. Granulomas (collections of epithelioid histiocytes) (Figure 11.20a) raise the possibility of mycobacterial infection (Figure 11.20b), fungal 's disease or a reaction infection, parasites, sarcoidosis, Crohn to foreign material, among numerous other possible causes. s may reflect parasitic infection or Eosinophils in large number allergy . Interpretation depends heavily on the site and clinical setting.

Figure 11.18 An acute inflammatory process characterised by numerous neutrophils. Note the typical multilobated nuclei (arrows). Figure 11.19 Oesophageal mucosa infiltrated by numerous eosinophils with bright red cytoplasm, many of which are forming clusters. Eosinophils may reflect allergy, parasitic infection or a wide variety of other causes. In this example, the clinicopathological diagnosis was eosinophilic oesophagitis.

Microscopic features of inflammation

Acute inflammation is characterised histologically by neutrophils (polymorphonuclear leukocytes), erosion or ulceration (Figure 11.18) and chronic inflammation by lymphocytes and plasma cells. Other inflammatory cells include eosinophils (Figure 11.19) , mast cells and histiocytes. Granulomas (collections of epithelioid histiocytes) (Figure 11.20a) raise the possibility of mycobacterial infection (Figure 11.20b), fungal 's disease or a reaction infection, parasites, sarcoidosis, Crohn to foreign material, among numerous other possible causes. s may reflect parasitic infection or Eosinophils in large number allergy . Interpretation depends heavily on the site and clinical setting.

Figure 11.18 An acute inflammatory process characterised by numerous neutrophils. Note the typical multilobated nuclei (arrows). Figure 11.19 Oesophageal mucosa infiltrated by numerous eosinophils with bright red cytoplasm, many of which are forming clusters. Eosinophils may reflect allergy, parasitic infection or a wide variety of other causes. In this example, the clinicopathological diagnosis was eosinophilic oesophagitis.

Revision #1

Created 2025-12-31 15:08:26 UTC by Omar Ayman

Updated 2025-12-31 15:08:26 UTC by Omar Ayman