

Minor salivary glands

Minor salivary glands

- Minor salivary glands appear at about the 12th week of gestation, forming directly from upper respiratory ectoderm. They develop as individual units with simple tubuloacinar systems. The minor salivary glands lack a distinct capsule and merge with the surrounding connective tissue in the submucosal region. They are widely distributed in the head and neck region and are mainly located (70-90%) in the oral cavity and oropharynx, including the palate, the tongue, the lips, the buccal mucosa and the retromolar trigone. The other sites are the nose, paranasal sinuses, pharynx and the larynx. They contribute 8-10% of the saliva and play a major role in saliva production during sleep. - Minor salivary glands
- Minor salivary glands appear at about the 12th week of gestation, forming directly from upper respiratory ectoderm. They develop as individual units with simple tubuloacinar systems. The minor salivary glands lack a distinct capsule and merge with the surrounding connective tissue in the submucosal region. They are widely distributed in the head and neck region and are mainly located (70-90%) in the oral cavity and oropharynx, including the palate, the tongue, the lips, the buccal mucosa and the retromolar trigone. The other sites are the nose, paranasal sinuses, pharynx and the larynx. They contribute 8-10% of the saliva and play a major role in saliva production during sleep. -

Revision #1

Created 2025-12-31 15:20:36 UTC by Omar Ayman

Updated 2025-12-31 15:20:36 UTC by Omar Ayman