

Nodes

Nodes

- Accurate assessment of nodal involvement remains a challenge for imaging. Most imaging techniques rely purely on size criteria to demonstrate lymph node involvement, with no possibility of identifying micrometastases in normal-sized nodes. A size criterion of 8–10 mm is often adopted, but it is not usually possible to distinguish benign reactive nodes from infiltrated nodes. This is a particular problem in patients with intrathoracic neoplasms, in whom enlarged benign reactive mediastinal nodes are common. The echo characteristics of nodes at endoscopic ultrasound have been used in many centres to increase the accuracy of nodal staging, and nodal sampling is possible via either mediastinoscopy or transoesophageal biopsy under endoscopic ultrasound control. PET/CT is of increasing use in detecting nodal metastases from a wide range of malignancies, with the capacity to co-register the area of increased FDG uptake with a precise anatomical location. Novel MRI contrast agents may help in the identification of non-enlarged tumour-infiltrated nodes.

Figure 8.41 (a) Endoscopic ultrasound in gastric cancer. The hypoechoic tumour (arrows) is infiltrating the layered structure of the gastric wall and extending out beyond the serosa. (b) Computed tomography scan demonstrates thickening and enhancement of the gastric wall in the same area (arrows). The stomach is distended with water to provide low-density contrast.

Nodes

- Accurate assessment of nodal involvement remains a challenge for imaging. Most imaging techniques rely purely on size criteria to demonstrate lymph node involvement, with no possibility of identifying micrometastases in normal-sized nodes. A size criterion of 8–10 mm is often adopted, but it is not usually possible to distinguish benign reactive nodes from infiltrated nodes. This is a particular problem in patients with intrathoracic neoplasms, in whom enlarged benign reactive mediastinal nodes are common. The echo characteristics of nodes at endoscopic ultrasound have been used in many centres to increase the accuracy of nodal staging, and nodal sampling is possible via either mediastinoscopy or transoesophageal biopsy under endoscopic ultrasound control. PET/CT is of increasing use in detecting nodal metastases from a wide range of malignancies, with the capacity to co-register the area of increased FDG uptake with a precise anatomical location. Novel MRI contrast agents may help in the identification of non-enlarged tumour-infiltrated nodes.

Figure 8.41 (a) Endoscopic ultrasound in gastric cancer. The hypoechoic tumour (arrows) is infiltrating the layered structure of the gastric wall and extending out beyond the serosa. (b) Computed tomography scan demonstrates thickening and enhancement of the gastric wall in the

same area (arrows). The stomach is distended with water to provide low-density contrast.

Nodes

- Accurate assessment of nodal involvement remains a challenge for imaging. Most imaging techniques rely purely on size criteria to demonstrate lymph node involvement, with no possibility of identifying micrometastases in normal-sized nodes. A size criterion of 8–10 mm is often adopted, but it is not usually possible to distinguish benign reactive nodes from infiltrated nodes. This is a particular problem in patients with intrathoracic neoplasms, in whom enlarged benign reactive mediastinal nodes are common. The echo characteristics of nodes at endoscopic ultrasound have been used in many centres to increase the accuracy of nodal staging, and nodal sampling is possible via either mediastinoscopy or transoesophageal biopsy under endoscopic ultrasound control. PET/CT is of increasing use in detecting nodal metastases from a wide range of malignancies, with the capacity to co-register the area of increased FDG uptake with a precise anatomical location. Novel MRI contrast agents may help in the identification of non-enlarged tumour-infiltrated nodes.

Figure 8.41 (a) Endoscopic ultrasound in gastric cancer. The hypoechoic tumour (arrows) is infiltrating the layered structure of the gastric wall and extending out beyond the serosa. (b) Computed tomography scan demonstrates thickening and enhancement of the gastric wall in the same area (arrows). The stomach is distended with water to provide low-density contrast.

Revision #1

Created 2025-12-31 15:28:52 UTC by Omar Ayman

Updated 2025-12-31 15:28:52 UTC by Omar Ayman