

Non-specific abdominal pain

Non-specific abdominal pain

The clinical features of NSAP are similar to acute appendicitis, but the pain is poorly localised, not aggravated by movement and not accompanied by guarding. The site and severity of maximum tenderness vary on repeated examinations. Symptoms are typically self-limiting. The aetiology is obscure, but viral infections are hypothesised if there is lymphadenopathy. In some children, recurrent abdominal pain can be organic; in others, it is eventually attributed to an underlying psychosocial problem. Non-specific abdominal pain

The clinical features of NSAP are similar to acute appendicitis, but the pain is poorly localised, not aggravated by movement and not accompanied by guarding. The site and severity of maximum tenderness vary on repeated examinations. Symptoms are typically self-limiting. The aetiology is obscure, but viral infections are hypothesised if there is lymphadenopathy. In some children, recurrent abdominal pain can be organic; in others, it is eventually attributed to an underlying psychosocial problem. Non-specific abdominal pain

The clinical features of NSAP are similar to acute appendicitis, but the pain is poorly localised, not aggravated by movement and not accompanied by guarding. The site and severity of maximum tenderness vary on repeated examinations. Symptoms are typically self-limiting. The aetiology is obscure, but viral infections are hypothesised if there is lymphadenopathy. In some children, recurrent abdominal pain can be organic; in others, it is eventually attributed to an underlying psychosocial problem.

Revision #1

Created 2025-12-31 15:18:29 UTC by Omar Ayman

Updated 2025-12-31 15:18:29 UTC by Omar Ayman