

Palpable undescended testes

Palpable undescended testes

Palpable UDTs require an orchidopexy at between 6 and 12 months. The canal is opened through an external oblique incision. The testis is mobilised on its vas and vessels, and the gubernaculum is usually divided. Any peritoneal outpocketing is ligated and divided at the internal ring, where dissection - adds length to the cord. The testis is placed in a subdartos scrotal pouch. Early orchidopexy , placing the testis in a cooler environment, improves spermatogenesis and may reduce the risk of testicular malignancy . Palpable undescended testes

Palpable UDTs require an orchidopexy at between 6 and 12 months. The canal is opened through an external oblique incision. The testis is mobilised on its vas and vessels, and the gubernaculum is usually divided. Any peritoneal outpocketing is ligated and divided at the internal ring, where dissection - adds length to the cord. The testis is placed in a subdartos scrotal pouch. Early orchidopexy , placing the testis in a cooler environment, improves spermatogenesis and may reduce the risk of testicular malignancy . Palpable undescended testes

Palpable UDTs require an orchidopexy at between 6 and 12 months. The canal is opened through an external oblique incision. The testis is mobilised on its vas and vessels, and the gubernaculum is usually divided. Any peritoneal outpocketing is ligated and divided at the internal ring, where dissection - adds length to the cord. The testis is placed in a subdartos scrotal pouch. Early orchidopexy , placing the testis in a cooler environment, improves spermatogenesis and may reduce the risk of testicular malignancy .

Revision #1

Created 2025-12-31 15:18:31 UTC by Omar Ayman

Updated 2025-12-31 15:18:31 UTC by Omar Ayman