

Polymerase chain reaction

Polymerase chain reaction

The polymerase chain reaction (PCR) amplifies DNA, yielding millions of copies from a single copy of a selected target. Amplification of RNA is also possible, using the technique of reverse transcriptase PCR (RT-PCR). It is worth noting that real-time PCR (RT-PCR) is a different method, typically used for quantification, with a very similar abbreviation. PCR is fast and safe and can be performed on homogenised fresh or formalin-fixed tissue. PCR-based methods have numerous applications in oncology (see Detection of clinically relevant abnormalities in genes), including mutational analysis (Figure 11.29), testing for clonality , detection of fusion transcripts resulting from cytogenetic changes, detection of amplifications, demonstration of MSI and detection of gene hypermethylation. PCR-based methods can also detect microorganisms in tissue but this is not a common application because of the risk of false positives. Polymerase chain reaction

The polymerase chain reaction (PCR) amplifies DNA, yielding millions of copies from a single copy of a selected target. Amplification of RNA is also possible, using the technique of reverse transcriptase PCR (RT-PCR). It is worth noting that real-time PCR (RT-PCR) is a different method, typically used for quantification, with a very similar abbreviation. PCR is fast and safe and can be performed on homogenised fresh or formalin-fixed tissue. PCR-based methods have numerous applications in oncology (see Detection of clinically relevant abnormalities in genes), including mutational analysis (Figure 11.29), testing for clonality , detection of fusion transcripts resulting from cytogenetic changes, detection of amplifications, demonstration of MSI and detection of gene hypermethylation. PCR-based methods can also detect microorganisms in tissue but this is not a common application because of the risk of false positives. Polymerase chain reaction

The polymerase chain reaction (PCR) amplifies DNA, yielding millions of copies from a single copy of a selected target. Amplification of RNA is also possible, using the technique of reverse transcriptase PCR (RT-PCR). It is worth noting that real-time PCR (RT-PCR) is a different method, typically used for quantification, with a very similar abbreviation. PCR is fast and safe and can be performed on homogenised fresh or formalin-fixed tissue. PCR-based methods have numerous applications in oncology (see Detection of clinically relevant abnormalities in genes), including mutational analysis (Figure 11.29), testing for clonality , detection of fusion transcripts resulting from cytogenetic changes, detection of amplifications, demonstration of MSI and detection of gene hypermethylation. PCR-based methods can also detect microorganisms in tissue but this is not a common application because of the risk of false positives.

Revision #1

Created 2025-12-31 15:08:29 UTC by Omar Ayman

Updated 2025-12-31 15:08:29 UTC by Omar Ayman