

# Rectum

## Rectum

Approximately 5% of colon injuries involve the rectum. These are generally from a penetrating injury , although occasionally the rectum may be damaged following fracture of the pelvis. Digital rectal examination will reveal the presence of blood, which is evidence of intestinal or rectal injury . These injuries are often associated with bladder and proximal urethral injury . With intraperitoneal injuries, the rectum is managed as for colonic injuries. Full-thickness extraperitoneal rectal injuries can be managed with primary repair and drainage depending on the type of injury , i.e. suitable for knife wounds but not ballistic trauma. Where there is extensive tissue loss, this should be managed with either a diverting end-colostomy and closure of the distal end (Hartmann's procedure) or a loop colostomy . Presacral drainage is no longer used. Rectum

Approximately 5% of colon injuries involve the rectum. These are generally from a penetrating injury , although occasionally the rectum may be damaged following fracture of the pelvis. Digital rectal examination will reveal the presence of blood, which is evidence of intestinal or rectal injury . These injuries are often associated with bladder and proximal urethral injury . With intraperitoneal injuries, the rectum is managed as for colonic injuries. Full-thickness extraperitoneal rectal injuries can be managed with primary repair and drainage depending on the type of injury , i.e. suitable for knife wounds but not ballistic trauma. Where there is extensive tissue loss, this should be managed with either a diverting end-colostomy and closure of the distal end (Hartmann's procedure) or a loop colostomy . Presacral drainage is no longer used. Rectum

Approximately 5% of colon injuries involve the rectum. These are generally from a penetrating injury , although occasionally the rectum may be damaged following fracture of the pelvis. Digital rectal examination will reveal the presence of blood, which is evidence of intestinal or rectal injury . These injuries are often associated with bladder and proximal urethral injury . With intraperitoneal injuries, the rectum is managed as for colonic injuries. Full-thickness extraperitoneal rectal injuries can be managed with primary repair and drainage depending on the type of injury , i.e. suitable for knife wounds but not ballistic trauma. Where there is extensive tissue loss, this should be managed with either a diverting end-colostomy and closure of the distal end (Hartmann's procedure) or a loop colostomy . Presacral drainage is no longer used.

---

Revision #1

Created 2025-12-31 15:12:36 UTC by Omar Ayman

Updated 2025-12-31 15:12:36 UTC by Omar Ayman