

# RECURRENT HYPERPARATHYROIDISM

## RECURRENT HYPERPARATHYROIDISM

Recurrent hyperparathyroidism is defined as hypercalcaemia occurring 6 months after surgery but with an intervening period of normocalcaemia. Common causes include missed pathology at the first operation; hyperplasia in remaining or autotransplanted tissue; parathyromatosis; or, very rarely, the development of a second parathyroid adenoma. Parathyromatosis refers to disseminated parathyroid tissue within the rupture of the parathyroid gland during the primary surgery. A definitive indication for surgical intervention must be present prior to embarking on localisation studies. Surgical intervention will be guided by the radiological imaging. Complication rates of recurrent laryngeal nerve damage and permanent hypocalcaemia are higher in reoperative surgery.

Recurrent hyperparathyroidism is defined as hypercalcaemia occurring 6 months after surgery but with an intervening period of normocalcaemia. Common causes include missed pathology at the first operation; hyperplasia in remaining or autotransplanted tissue; parathyromatosis; or, very rarely, the development of a second parathyroid adenoma. Parathyromatosis refers to disseminated parathyroid tissue within the rupture of the parathyroid gland during the primary surgery. A definitive indication for surgical intervention must be present prior to embarking on localisation studies. Surgical intervention will be guided by the radiological imaging. Complication rates of recurrent laryngeal nerve damage and permanent hypocalcaemia are higher in reoperative surgery.

---

Revision #1

Created 2025-12-31 15:21:06 UTC by Omar Ayman

Updated 2025-12-31 15:21:06 UTC by Omar Ayman