

REFERENCES

REFERENCES

1 Coelho KR. Challenges of the oral cancer burden in India. J Cancer Epidemiol 2012; 2012 : 701932. 2 Warnakulasuriya S, Ariyawardana A. Malignant transformation of oral leukoplakia: a systematic review of observational studies. J Oral Pathol Med 2016; 45 (3): 155-66. 3 D'Cruz AK, Vaish R, Kapre N et al . Elective versus therapeutic neck dissection in node-negative oral cancer. N Engl J Med 2015; 373 (6): 521-9. 4 Schilling C, Stoeckli SJ, Haerle SK et al . Sentinel European Node Trial (SENT): 3-year results of sentinel node biopsy in oral cancer. Eur J Cancer 2015; 51 (18): 2777-84. - 5 Hutchison IL, Ridout F , Cheung SMY et al . Nationwide randomised trial evaluating elective neck dissection for early stage oral cancer (SEND study) with meta-analysis and concurrent real-world cohort. Br J Cancer 2019; 121 (10): 827-36. - 6 Lindberg R. Distribution of cervical lymph node metastases from squamous cell carcinoma of the upper respiratory and digestive tracts. Cancer 1972; 29 (6): 1446-9. 7 Shah JP . Patterns of cervical lymph node metastasis from squamous carcinomas of the upper aerodigestive tract. Am J Surg 1990; 160 (4): 405-9. 8 Expert Panel on Radiation Oncology-Head and Neck; Salama JK, Saba N, Quon H et al . ACR appropriateness criteria® adjuvant ther - apy for resected squamous cell carcinoma of the head and neck. Oral Oncol 2011; 47 (7): 554-9. REFERENCES

1 Coelho KR. Challenges of the oral cancer burden in India. J Cancer Epidemiol 2012; 2012 : 701932. 2 Warnakulasuriya S, Ariyawardana A. Malignant transformation of oral leukoplakia: a systematic review of observational studies. J Oral Pathol Med 2016; 45 (3): 155-66. 3 D'Cruz AK, Vaish R, Kapre N et al . Elective versus therapeutic neck dissection in node-negative oral cancer. N Engl J Med 2015; 373 (6): 521-9. 4 Schilling C, Stoeckli SJ, Haerle SK et al . Sentinel European Node Trial (SENT): 3-year results of sentinel node biopsy in oral cancer. Eur J Cancer 2015; 51 (18): 2777-84. - 5 Hutchison IL, Ridout F , Cheung SMY et al . Nationwide randomised trial evaluating elective neck dissection for early stage oral cancer (SEND study) with meta-analysis and concurrent real-world cohort. Br J Cancer 2019; 121 (10): 827-36. - 6 Lindberg R. Distribution of cervical lymph node metastases from squamous cell carcinoma of the upper respiratory and digestive tracts. Cancer 1972; 29 (6): 1446-9. 7 Shah JP . Patterns of cervical lymph node metastasis from squamous carcinomas of the upper aerodigestive tract. Am J Surg 1990; 160 (4): 405-9. 8 Expert Panel on Radiation Oncology-Head and Neck; Salama JK, Saba N, Quon H et al . ACR appropriateness criteria® adjuvant ther - apy for resected squamous cell carcinoma of the head and neck. Oral Oncol 2011; 47 (7): 554-9.

Revision #1

Created 2025-12-31 15:20:25 UTC by Omar Ayman

Updated 2025-12-31 15:20:25 UTC by Omar Ayman