

Regional anaesthesia

Regional anaesthesia

Regional anaesthesia involves central neuraxial or peripheral nerve or plexus blocks using local anaesthetic drugs. It has a clear advantage when general anaesthesia carries a higher risk of morbidity and mortality, such as in patients with debilitating respiratory and cardiovascular disease and obstetric cases. It also provides excellent pain relief in the postoperative period, reducing the need for analgesics such as opioids. As with general anaesthesia, venous access should be established and vital parameters should be monitored during regional anaesthesia. Localising nerves using anatomical landmarks and eliciting paraesthesia alone carries a high risk of nerve damage and intravascular injection and has a lower success rate. The use of nerve stimulators to localise nerves improves the success rate and reduces risks. Ultrasound-guided regional anaesthesia allows the visualisation of nerves and the spread of local anaesthetics, enabling the use of a smaller dose of local anaesthetic agents with improved success rates and safety. Summary box 23.8 Types of anaesthesia

General anaesthesia may be more acceptable to patients Regional anaesthesia has major advantages in obstetrics and patients with respiratory compromise Local blocks have been transformed by nerve stimulators and ultrasound guidance All require full resuscitation and monitoring equipment to be available

Regional anaesthesia

Regional anaesthesia involves central neuraxial or peripheral nerve or plexus blocks using local anaesthetic drugs. It has a clear advantage when general anaesthesia carries a higher risk of morbidity and mortality, such as in patients with debilitating respiratory and cardiovascular disease and obstetric cases. It also provides excellent pain relief in the postoperative period, reducing the need for analgesics such as opioids. As with general anaesthesia, venous access should be established and vital parameters should be monitored during regional anaesthesia. Localising nerves using anatomical landmarks and eliciting paraesthesia alone carries a high risk of nerve damage and intravascular injection and has a lower success rate. The use of nerve stimulators to localise nerves improves the success rate and reduces risks. Ultrasound-guided regional anaesthesia allows the visualisation of nerves and the spread of local anaesthetics, enabling the use of a smaller dose of local anaesthetic agents with improved success rates and safety. Summary box 23.8 Types of anaesthesia

General anaesthesia may be more acceptable to patients Regional anaesthesia has major advantages in obstetrics and patients with respiratory compromise Local blocks have been transformed by nerve stimulators and ultrasound guidance All require full resuscitation and monitoring equipment to be available

Regional anaesthesia

Regional anaesthesia involves central neuraxial or peripheral nerve or plexus blocks using local anaesthetic drugs. It has a clear advantage when general anaesthesia carries a higher risk of morbidity and mortality, such as in patients with debilitating respiratory and cardiovascular disease and obstetric cases. It also provides excellent pain relief in the postoperative period, reducing the need for analgesics such as opioids. As with general anaesthesia, venous access should be established and vital parameters should be monitored during regional anaesthesia. Localising nerves using anatomical landmarks and eliciting paraesthesia alone carries a high risk of nerve damage and intravascular injection and has a lower success rate. The use of nerve stimulators to localise nerves improves the success rate and reduces risks. Ultrasound-guided regional anaesthesia allows the visualisation of nerves and the spread of local anaesthetics, enabling the use of a smaller dose of local anaesthetic agents with improved success rates and safety. Summary box 23.8 Types of anaesthesia /uni25CF /uni25CF /uni25CF /uni25CF

General anaesthesia may be more acceptable to patients Regional anaesthesia has major advantages in obstetrics and patients with respiratory compromise Local blocks have been transformed by nerve stimulators and ultrasound guidance All require full resuscitation and monitoring equipment to be available

Revision #1

Created 2025-12-31 15:11:13 UTC by Omar Ayman

Updated 2025-12-31 15:11:13 UTC by Omar Ayman