

# Revision knee replacement

## Revision knee replacement

TKRs have excellent long-term survival, with 82% lasting over 20 years and over 95% lasting 10–15 years. In the first few years, revision surgery is most likely to be due to infection, and non-infective causes, such as instability and malalignment. Beyond 10 years, revision is usually a result of aseptic loosening of components, which is often a result of wear of the plastic insert and polyethylene-induced osteolysis. With any failed joint replacement, infection should be excluded as this can compromise the result of surgery and may require a different surgical strategy, e.g. a two-stage rather than a single-stage procedure. Revision arthroplasty is technically more challenging, requires more complex and expensive implants and has a higher complication rate. When done for the correct indication, results are very good. As with a primary TKR, the goal remains to provide a well-aligned, stable and pain-free knee ( Figure 40.6 ).

(b) (c) (d) Figure 40.6 Radiographs of a malaligned knee (a, b) and a well-aligned revised knee (c, d) .

## Revision knee replacement

TKRs have excellent long-term survival, with 82% lasting over 20 years and over 95% lasting 10–15 years. In the first few years, revision surgery is most likely to be due to infection, and non-infective causes, such as instability and malalignment. Beyond 10 years, revision is usually a result of aseptic loosening of components, which is often a result of wear of the plastic insert and polyethylene-induced osteolysis. With any failed joint replacement, infection should be excluded as this can compromise the result of surgery and may require a different surgical strategy, e.g. a two-stage rather than a single-stage procedure. Revision arthroplasty is technically more challenging, requires more complex and expensive implants and has a higher complication rate. When done for the correct indication, results are very good. As with a primary TKR, the goal remains to provide a well-aligned, stable and pain-free knee ( Figure 40.6 ).

(b) (c) (d) Figure 40.6 Radiographs of a malaligned knee (a, b) and a well-aligned revised knee (c, d) .

## Revision knee replacement

TKRs have excellent long-term survival, with 82% lasting over 20 years and over 95% lasting 10–15 years. In the first few years, revision surgery is most likely to be due to infection, and non-infective causes, such as instability and malalignment. Beyond 10 years, revision is usually a result of aseptic loosening of components, which is often a result of wear of the plastic insert and polyethylene-induced

osteolysis. With any failed joint replacement, infection should be excluded as this can compromise the result of surgery and may require a different surgical strategy, e.g. a two-stage rather than a single-stage procedure. Revision arthroplasty is technically more challenging, requires more complex and expensive implants and has a higher complication rate. When done for the correct indication, results are very good. As with a primary TKR, the goal remains to provide a well-aligned, stable and pain-free knee ( Figure 40.6 ).

(b) (c) (d) Figure 40.6 Radiographs of a malaligned knee (a, b) and a well-aligned revised knee (c, d) .

---

Revision #1

Created 2025-12-31 15:16:04 UTC by Omar Ayman

Updated 2025-12-31 15:16:04 UTC by Omar Ayman