

Sacrococcygeal teratoma

Sacrococcygeal teratoma

These germ cell tumours arise from the coccyx and are usually diagnosed antenatally . They may have internal and external components or both (Figure 18.20). Most are benign mature teratomas, but some contain immature embryonic elements. Complete excision is usually achieved in the prone position the pelvic floor. If the intra-abdominal component is large or vascular, the median sacral artery can be ligated through the abdomen. Bladder and bowel function are assessed following the pelvic floor repair. α -Fetoprotein levels are measured to detect recurrence.

Figure 18.20 Sacrococcygeal teratoma.

Sacrococcygeal teratoma

These germ cell tumours arise from the coccyx and are usually diagnosed antenatally . They may have internal and external components or both (Figure 18.20). Most are benign mature teratomas, but some contain immature embryonic elements. Complete excision is usually achieved in the prone position the pelvic floor. If the intra-abdominal component is large or vascular, the median sacral artery can be ligated through the abdomen. Bladder and bowel function are assessed following the pelvic floor repair. α -Fetoprotein levels are measured to detect recurrence.

Figure 18.20 Sacrococcygeal teratoma.

Sacrococcygeal teratoma

These germ cell tumours arise from the coccyx and are usually diagnosed antenatally . They may have internal and external components or both (Figure 18.20). Most are benign mature teratomas, but some contain immature embryonic elements. Complete excision is usually achieved in the prone position the pelvic floor. If the intra-abdominal component is large or vascular, the median sacral artery can be ligated through the abdomen. Bladder and bowel function are assessed following the pelvic floor repair. α -Fetoprotein levels are measured to detect recurrence.

Figure 18.20 Sacrococcygeal teratoma.

Revision #1

Created 2025-12-31 15:09:58 UTC by Omar Ayman

Updated 2025-12-31 15:09:58 UTC by Omar Ayman