

SECONDARY SURVEY

SECONDARY SURVEY

All severely injured patients require a detailed top-to-toe examination after life-threatening injuries have been identified and managed during the primary survey. Patients may be intubated and unresponsive at this point, limiting the accuracy of clinical examination. Such patients should have a 'tertiary survey' when extubated and alert to identify any missed 'minor' injuries, such as a scaphoid fracture in the wrist or a rotator cuff tear in the shoulder. These injuries have the potential to cause significant long-term disability. It is essential that the findings of the primary, secondary and tertiary surveys are clearly recorded in the patient's case notes. SECONDARY SURVEY

All severely injured patients require a detailed top-to-toe examination after life-threatening injuries have been identified and managed during the primary survey. Patients may be intubated and unresponsive at this point, limiting the accuracy of clinical examination. Such patients should have a 'tertiary survey' when extubated and alert to identify any missed 'minor' injuries, such as a scaphoid fracture in the wrist or a rotator cuff tear in the shoulder. These injuries have the potential to cause significant long-term disability. It is essential that the findings of the primary, secondary and tertiary surveys are clearly recorded in the patient's case notes.

Revision #1

Created 2025-12-31 15:12:06 UTC by Omar Ayman

Updated 2025-12-31 15:12:06 UTC by Omar Ayman