

Sialorrhoea

Sialorrhoea

Excess salivation is rarely symptomatic in healthy individuals as it is swallowed spontaneously . Certain drugs and oral infection produce a transient increase in salivary flow; however, uncontrolled drooling is usually seen in the presence of normal salivary production in children with mental and physical disability , most notably cerebral palsy . Sialorrhoea can be managed with antisialogogues or with intraparenchymal injection of botulinum toxin injection. Most resting salivary flow arises from the submandibular glands and surgery should focus on either bilateral excision or repositioning of the salivary duct to control sialorrhoea. Sialorrhoea

Excess salivation is rarely symptomatic in healthy individuals as it is swallowed spontaneously . Certain drugs and oral infection produce a transient increase in salivary flow; however, uncontrolled drooling is usually seen in the presence of normal salivary production in children with mental and physical disability , most notably cerebral palsy . Sialorrhoea can be managed with antisialogogues or with intraparenchymal injection of botulinum toxin injection. Most resting salivary flow arises from the submandibular glands and surgery should focus on either bilateral excision or repositioning of the salivary duct to control sialorrhoea.

Revision #1

Created 2025-12-31 15:20:39 UTC by Omar Ayman

Updated 2025-12-31 15:20:39 UTC by Omar Ayman