

Spinal examination

Spinal examination

The overlying skin should be inspected (e.g. for possible penetrating wounds) and the entire spine must be palpated. A formal spinal log roll must be performed to achieve this (Figure 30.12). Significant swelling, tenderness, palpable steps or gaps suggest a spinal injury . A rectal examination should be undertaken to assess anal tone and perianal sensation (see Neurological examination). Seatbelt marks on the abdomen and chest must be noted, as these suggest a high-energy accident. Spinal examination

The overlying skin should be inspected (e.g. for possible penetrating wounds) and the entire spine must be palpated. A formal spinal log roll must be performed to achieve this (Figure 30.12). Significant swelling, tenderness, palpable steps or gaps suggest a spinal injury . A rectal examination should be undertaken to assess anal tone and perianal sensation (see Neurological examination). Seatbelt marks on the abdomen and chest must be noted, as these suggest a high-energy accident. Spinal examination

The overlying skin should be inspected (e.g. for possible penetrating wounds) and the entire spine must be palpated. A formal spinal log roll must be performed to achieve this (Figure 30.12). Significant swelling, tenderness, palpable steps or gaps suggest a spinal injury . A rectal examination should be undertaken to assess anal tone and perianal sensation (see Neurological examination). Seatbelt marks on the abdomen and chest must be noted, as these suggest a high-energy accident.

Revision #1

Created 2025-12-31 15:13:08 UTC by Omar Ayman

Updated 2025-12-31 15:13:08 UTC by Omar Ayman