

Syringomyelia

Syringomyelia

Patients may present with sensory disturbance, weakness of the hands, loss of pain and temperature sensation, asymmetrical abdominal reflexes or progressive kyphoscoliosis. It is associated with Arnold–Chiari malformation and spinal cord tumours. Where syringomyelia is associated with an Arnold–Chiari malformation and scoliosis, a posterior cranial fossa decompression should be carried out first to resolve the syringomyelia. The scoliosis may then be corrected at a later date. Syringomyelia

Patients may present with sensory disturbance, weakness of the hands, loss of pain and temperature sensation, asymmetrical abdominal reflexes or progressive kyphoscoliosis. It is associated with Arnold–Chiari malformation and spinal cord tumours. Where syringomyelia is associated with an Arnold–Chiari malformation and scoliosis, a posterior cranial fossa decompression should be carried out first to resolve the syringomyelia. The scoliosis may then be corrected at a later date. Syringomyelia

Patients may present with sensory disturbance, weakness of the hands, loss of pain and temperature sensation, asymmetrical abdominal reflexes or progressive kyphoscoliosis. It is associated with Arnold–Chiari malformation and spinal cord tumours. Where syringomyelia is associated with an Arnold–Chiari malformation and scoliosis, a posterior cranial fossa decompression should be carried out first to resolve the syringomyelia. The scoliosis may then be corrected at a later date.

Revision #1

Created 2025-12-31 15:15:21 UTC by Omar Ayman

Updated 2025-12-31 15:15:21 UTC by Omar Ayman