

# Tarsometatarsal (Lisfranc) joint injuries

## Tarsometatarsal (Lisfranc) joint injuries

- Injuries to the midfoot are associated with significant morbidity ranging from a midfoot sprain to complete rupture of the ligaments connecting the forefoot to the midfoot. Injury classically follows forced plantarflexion of the midfoot. An alternative mechanism of injury are crush injuries where the foot is forced flat by a heavy weight. Lisfranc's ligament connects the second metatarsal to the medial cuneiform. Poorly treated injuries to the midfoot lead to significant morbidity and, if suspected, a CT of the foot should be undertaken. Treatment options range from closed reduction and plaster cast application to open reduction and internal fixation. In severe cases primary tarsometatarsal fusion may be considered. Tarsometatarsal (Lisfranc) joint injuries
- Injuries to the midfoot are associated with significant morbidity ranging from a midfoot sprain to complete rupture of the ligaments connecting the forefoot to the midfoot. Injury classically follows forced plantarflexion of the midfoot. An alternative mechanism of injury are crush injuries where the foot is forced flat by a heavy weight. Lisfranc's ligament connects the second metatarsal to the medial cuneiform. Poorly treated injuries to the midfoot lead to significant morbidity and, if suspected, a CT of the foot should be undertaken. Treatment options range from closed reduction and plaster cast application to open reduction and internal fixation. In severe cases primary tarsometatarsal fusion may be considered. Tarsometatarsal (Lisfranc) joint injuries
- Injuries to the midfoot are associated with significant morbidity ranging from a midfoot sprain to complete rupture of the ligaments connecting the forefoot to the midfoot. Injury classically follows forced plantarflexion of the midfoot. An alternative mechanism of injury are crush injuries where the foot is forced flat by a heavy weight. Lisfranc's ligament connects the second metatarsal to the medial cuneiform. Poorly treated injuries to the midfoot lead to significant morbidity and, if suspected, a CT of the foot should be undertaken. Treatment options range from closed reduction and plaster cast application to open reduction and internal fixation. In severe cases primary tarsometatarsal fusion may be considered.

---

Revision #1

Created 2025-12-31 15:13:41 UTC by Omar Ayman

Updated 2025-12-31 15:13:41 UTC by Omar Ayman