

Tendon

Tendon

Although repair follows the normal pattern of wound healing, there are two main mechanisms whereby nutrients, cells and new vessels reach the severed tendon. Intrinsic healing consists of vincular blood flow and synovial diffusion. Extrinsic healing depends on the formation of fibrous adhesions between the tendon and the tendon sheath. Early active mobilisation following tendon repairs prevents adhesions limiting the range of motion and therefore promotes the more desired intrinsic healing. At the same time, tendon repairs must be protected by splintage to avoid rupture. Tendon

Although repair follows the normal pattern of wound healing, there are two main mechanisms whereby nutrients, cells and new vessels reach the severed tendon. Intrinsic healing consists of vincular blood flow and synovial diffusion. Extrinsic healing depends on the formation of fibrous adhesions between the tendon and the tendon sheath. Early active mobilisation following tendon repairs prevents adhesions limiting the range of motion and therefore promotes the more desired intrinsic healing. At the same time, tendon repairs must be protected by splintage to avoid rupture. Tendon

Although repair follows the normal pattern of wound healing, there are two main mechanisms whereby nutrients, cells and new vessels reach the severed tendon. Intrinsic healing consists of vincular blood flow and synovial diffusion. Extrinsic healing depends on the formation of fibrous adhesions between the tendon and the tendon sheath. Early active mobilisation following tendon repairs prevents adhesions limiting the range of motion and therefore promotes the more desired intrinsic healing. At the same time, tendon repairs must be protected by splintage to avoid rupture.

Revision #1

Created 2025-12-31 15:12:50 UTC by Omar Ayman

Updated 2025-12-31 15:12:50 UTC by Omar Ayman