

TYPHOID Introduction

TYPHOID Introduction

Typhoid fever is caused by *Salmonella Typhi*, also called the typhoid bacillus, a Gram-negative organism. Like most infections occurring in the tropics, the organism gains entry into the human gastrointestinal tract as a result of poor hygiene and inadequate sanitation. It is a disease normally managed by physicians, but the surgeon may be called upon to treat the patient with typhoid fever because of perforation of a typhoid ulcer. TYPHOID Introduction

Typhoid fever is caused by *Salmonella Typhi*, also called the typhoid bacillus, a Gram-negative organism. Like most infections occurring in the tropics, the organism gains entry into the human gastrointestinal tract as a result of poor hygiene and inadequate sanitation. It is a disease normally managed by physicians, but the surgeon may be called upon to treat the patient with typhoid fever because of perforation of a typhoid ulcer. TYPHOID Introduction

Typhoid fever is caused by *Salmonella Typhi*, also called the typhoid bacillus, a Gram-negative organism. Like most infections occurring in the tropics, the organism gains entry into the human gastrointestinal tract as a result of poor hygiene and inadequate sanitation. It is a disease normally managed by physicians, but the surgeon may be called upon to treat the patient with typhoid fever because of perforation of a typhoid ulcer.

Revision #1

Created 2025-12-31 15:22:39 UTC by Omar Ayman

Updated 2025-12-31 15:22:39 UTC by Omar Ayman