

Urinary catheter

Urinary catheter

The requirement for a urinary catheter depends on the operation. In shorter (<4 hours) minimal access procedures a urinary catheter is not usually required. If a urinary catheter has been placed in the bladder during an operation with likely short stay, it can be removed before the patient regains consciousness if the procedure has been uneventful. Postoperatively it is important to check that the patient has been able to pass urine and empty their bladder without difficulty. When there is uncertainty point-of-care bladder scanning can assess residual bladder volume.

Urinary catheter

The requirement for a urinary catheter depends on the operation. In shorter (<4 hours) minimal access procedures a urinary catheter is not usually required. If a urinary catheter has been placed in the bladder during an operation with likely short stay, it can be removed before the patient regains consciousness if the procedure has been uneventful. Postoperatively it is important to check that the patient has been able to pass urine and empty their bladder without difficulty. When there is uncertainty point-of-care bladder scanning can assess residual bladder volume.

Urinary catheter

The requirement for a urinary catheter depends on the operation. In shorter (<4 hours) minimal access procedures a urinary catheter is not usually required. If a urinary catheter has been placed in the bladder during an operation with likely short stay, it can be removed before the patient regains consciousness if the procedure has been uneventful. Postoperatively it is important to check that the patient has been able to pass urine and empty their bladder without difficulty. When there is uncertainty point-of-care bladder scanning can assess residual bladder volume.

Revision #1

Created 2025-12-31 15:08:09 UTC by Omar Ayman

Updated 2025-12-31 15:08:09 UTC by Omar Ayman