

# 54 - References

## References

### Prescribing in children and adolescents CHAPTER 5 References

1. Kendrick JG, et al. Pharmacologic management of agitation and aggression in a pediatric emergency department - a retrospective cohort study. *J Pediatric Pharmacol Therapeutics* 2018; 23:455-459.
2. Wolpert KH, et al. Behavioral management of children with autism in the emergency department. *Pediatr Emerg Care* 2023; 39:45-50.
3. National Institute for Health and Care Excellence. Psychosis and schizophrenia in children and young people: recognition and management. Clinical Guidance [CG155]. 2013 (last updated October 2016, last checked October 2023); <https://www.nice.org.uk/guidance/cg155>.
4. Chang MY, et al. Drug-induced extrapyramidal symptoms at the pediatric emergency department. *Pediatr Emerg Care* 2020; 36:468-472.
5. National Institute for Health and Care Excellence. Violence and aggression: short-term management in mental health, health and community settings. NICE Guideline [NG10]. 2015 (last checked October 2023); <https://www.nice.org.uk/guidance/ng10>.
6. Rudolf F, et al. A retrospective review of antipsychotic medications administered to psychiatric patients in a tertiary care pediatric emergency department. *J Pediatric Pharmacol Therapeutics* 2019; 24:234-237.
7. Cole JB, et al. The use, safety, and efficacy of olanzapine in a level I pediatric trauma center emergency department over a 10-year period. *Pediatr Emerg Care* 2020; 36:70-76.
8. Breier A, et al. A double-blind, placebo-controlled dose-response comparison of intramuscular olanzapine and haloperidol in the treatment of acute agitation in schizophrenia. *Arch Gen Psychiatry* 2002; 59:441-448.
9. Lindborg SR, et al. Effects of intramuscular olanzapine vs. haloperidol and placebo on QTc intervals in acutely agitated patients. *Psychiatry Res* 2003; 119:113-123.
10. Powney MJ, et al. Haloperidol for psychosis-induced aggression or agitation (rapid tranquillisation). *Cochrane Database Syst Rev* 2012; 11:CD009377.
11. TREC Collaborative Group. Rapid tranquillisation for agitated patients in emergency psychiatric rooms: a randomised trial of midazolam versus haloperidol plus promethazine. *BMJ* 2003; 327:708-713.
12. Nobay F, et al. A prospective, double-blind, randomized trial of midazolam versus haloperidol versus lorazepam in the chemical restraint of violent and severely agitated patients. *Acad Emerg Med* 2004; 11:744-749.

13. Schwagmeier R, et al. Midazolam pharmacokinetics following intravenous and buccal administration. *Br J Clin Pharmacol* 1998; 46:203–206.
14. Taylor D, et al. Buccal midazolam for agitation on psychiatric intensive care wards. *Int J Psychiatry Clin Pract* 2008; 12:309–311.
15. Nunn K, et al. Medication table. In: Nunn KP, Dey C, eds. *The Clinician's Guide to Psychotropic Prescribing in Children and Adolescents*, 1st edn. Sydney: Glad Publishing 2003:383–452.
16. Khan SS, et al. A naturalistic evaluation of intramuscular ziprasidone versus intramuscular olanzapine for the management of acute agitation and aggression in children and adolescents. *J Child Adolesc Psychopharmacol* 2006; 16:671–677.
17. Staller JA. Intramuscular ziprasidone in youth: a retrospective chart review. *J Child Adolesc Psychopharmacol* 2004; 14:590–592.
18. Sanford M, et al. Intramuscular aripiprazole: a review of its use in the management of agitation in schizophrenia and bipolar I disorder. *CNS Drugs* 2008; 22:335–352.
19. National Institute for Health and Clinical Excellence. Aripiprazole for schizophrenia in people aged 15 to 17 years. Technology Appraisal [TA213]. 2011 (last checked October 2023); <https://www.nice.org.uk/guidance/ta213>.
20. Lesem MD, et al. Rapid acute treatment of agitation in individuals with schizophrenia: multicentre, randomised, placebo-controlled study of inhaled loxapine. *Br J Psychiatry* 2011; 198:51–58.

---

Revision #1

Created 2026-01-04 20:16:48 UTC by Omar Ayman

Updated 2026-01-04 20:16:48 UTC by Omar Ayman